

FREE QUASI-SYMMETRIC FUNCTIONS AND DESCENT ALGEBRAS FOR WREATH PRODUCTS, AND NONCOMMUTATIVE MULTI-SYMMETRIC FUNCTIONS

JEAN-CHRISTOPHE NOVELLI AND JEAN-YVES THIBON

ABSTRACT. We introduce analogs of the Hopf algebra of Free quasi-symmetric functions with bases labelled by colored permutations. When the color set is a semigroup, an internal product can be introduced. This leads to the construction of generalized descent algebras associated with wreath products $\Gamma \wr \mathfrak{S}_n$ and to the corresponding generalizations of quasi-symmetric functions. The associated Hopf algebras appear as natural analogs of McMahon's multisymmetric functions. As a consequence, we obtain an internal product on ordinary multi-symmetric functions. We extend these constructions to Hopf algebras of colored parking functions, colored non-crossing partitions and parking functions of type B .

CONTENTS

1. Introduction	2
2. Background and notations	3
2.1. Colored alphabets	4
2.2. Colored standardization	4
2.3. Colored shifted operations	4
3. Free quasi-symmetric functions of level ℓ	5
3.1. $\mathbf{FQSym}^{(\ell)}$ and $\mathbf{FQSym}^{(\Gamma)}$	5
3.2. Duality in $\mathbf{FQSym}^{(\ell)}$	6
3.3. Algebraic structure of $\mathbf{FQSym}^{(\ell)}$	8
3.4. Primitive elements of $\mathbf{FQSym}^{(\ell)}$	8
3.5. Dendriform structure of $\mathbf{FQSym}^{(\ell)}$	9
3.6. Internal product of $\mathbf{FQSym}^{(\ell)}$	9
4. Noncommutative symmetric functions of level ℓ	10
4.1. ℓ -partite numbers	10
4.2. The Hopf algebra $\mathbf{Sym}^{(\ell)}$	10
4.3. Algebraic structure of $\mathbf{Sym}^{(\ell)}$	11
4.4. Primitive elements of $\mathbf{Sym}^{(\ell)}$	12
4.5. Duality in $\mathbf{Sym}^{(\ell)}$	12
5. Quasi-symmetric functions of level ℓ	13
5.1. Cauchy formula of level ℓ	13
5.2. Poirier's Quasi-symmetric functions	13
5.3. Coproducts and alphabets	14

Date: June 24, 2008.

5.4. Generalized descent algebras	17
5.5. Ordinary multi-symmetric functions	18
6. The Mantaci-Reutenauer algebra	19
6.1. Monochromatic complete functions	19
6.2. Primitive elements of $\text{MR}^{(\ell)}$	20
6.3. Duality	21
7. Level ℓ parking quasi-symmetric functions	21
7.1. Usual parking functions	21
7.2. Colored parking functions	22
7.3. Duality	23
7.4. Algebraic structure of $\mathbf{PQSym}^{(\ell)}$	23
7.5. Primitive elements of $\mathbf{PQSym}^{(\ell)}$	24
7.6. Bidendriform and tridendriform structure	25
8. Type B algebras	26
8.1. Parking functions of type B	26
8.2. Non-crossing partitions of type B	26
9. Colored analogs of planar binary trees: $\mathbf{PBT}^{(\ell)}$	27
9.1. Definition of $\mathbf{PBT}^{(\ell)}$	27
9.2. Algebraic structure of $\mathbf{PBT}^{(\ell)}$	28
10. Examples	29
10.1. Multigraded coinvariants and colored Klyachko idempotents	29
10.2. A formula of Raney	32
References	33

1. INTRODUCTION

The Hopf algebra of Free Quasi-Symmetric Functions $\mathbf{FQSym}$ [4] is an algebra of noncommutative polynomials associated with the sequence $(\mathfrak{S}_n)_{n \geq 0}$ of all symmetric groups. It is connected by Hopf homomorphisms to several other important algebras associated with the same sequence of groups : Free symmetric functions (or coplactic algebra) $\mathbf{FSym}$ [36, 4], Non-commutative symmetric functions (or descent algebras) $\mathbf{Sym}$ [7], Quasi-Symmetric functions $QSym$ [9], Symmetric functions Sym , and also, Planar binary trees $\mathbf{PBT}$ [22, 12], Matrix quasi-symmetric functions $\mathbf{MQSym}$ [4, 11], Parking functions $\mathbf{PQSym}$ [19, 29], and so on.

Most of these Hopf algebras are endowed with an internal product, generalizing the one of ordinary symmetric functions. The basic example is provided by noncommutative symmetric functions, whose homogeneous components can be identified with the Solomon descent algebras of symmetric groups [7].

Symmetric groups are the Coxeter groups of type A , and there are descent algebras for other types as well. However, the direct sums of the descent algebras of types B or D are not Hopf algebras in any natural way. But there are Hopf algebras associated with wreath products $\mathbb{Z}_\ell \wr \mathfrak{S}_n$, the Mantaci-Reutenauer algebras [27], which admit internal products, and contain the Solomon algebras of type B for $\ell = 2$.

From the point of view of symmetric functions, $\text{MR}^{(\ell)}$, the Mantaci-Reutenauer algebra of level ℓ is the free product of ℓ copies of **Sym**. It is therefore the natural noncommutative analog of $(\text{Sym})^{\otimes \ell} \simeq \text{Sym}(X_0; \dots; X_{\ell-1})$, the algebra of symmetric functions in ℓ independent alphabets, which is also the Grothendieck ring of the tower of algebras $(\mathbb{C}[\mathbb{Z}_\ell \wr \mathfrak{S}_n])$. And indeed, it has been shown in [14] that $\text{MR}^{(\ell)}$ was the Grothendieck ring of projective modules over the 0-Ariki-Koike-Shoji algebras, a degeneracy of the Hecke algebras associated with $\mathbb{Z}_\ell \wr \mathfrak{S}_n$.

However, with ℓ independent alphabets, one can build a larger Hopf algebras. In the commutative case, it is the algebra of *multi-symmetric functions*, first introduced by McMahon [26], and briefly investigated by Gessel from a modern point of view in [10]. It is defined as follows. Setting $X_i = \{x_{i,j} | j = 1, \dots, n\}$, the multi-symmetric polynomials are the invariants of $\mathfrak{S}_n$ in $\mathbb{C}[X_0, \dots, X_{\ell-1}]$ for the diagonal action (by the automorphisms $\sigma(x_{i,j}) = x_{i,\sigma(j)}$). This is an algebra, which, as usual, acquires a Hopf algebra structure in the limit $n \rightarrow \infty$.

In the following, we will start with a level ℓ analogue of **FQSym**, whose bases are labelled by ℓ -colored permutations. Imitating the embedding of **Sym** in **FQSym**, we obtain a Hopf subalgebra of level ℓ called $\mathbf{Sym}^{(\ell)}$, which is a natural noncommutative analog of McMahon's algebra of multi-symmetric functions, and turns out to be dual to Poirier's quasi-symmetric functions. Its homogenous components can be endowed with an internal product, thus providing an analog of Solomon's descent algebras for wreath products, bigger than the Mantaci-Reutenauer algebras, and in which most useful properties such as the splitting formula remain valid. By commutative image, this yields an internal product on multi-symmetric functions.

The Mantaci-Reutenauer descent algebra $\text{MR}^{(\ell)}$ arises as a natural Hopf subalgebra of $\mathbf{Sym}^{(\ell)}$ and its dual is computed in a straightforward way by means of an appropriate Cauchy formula.

Finally, we introduce a Hopf algebra of colored parking functions, and use it to define Hopf algebras structures on parking functions and non-crossing partitions of type B .

The main results of this paper have been announced in the draft [32]. Since then, some of these results, in particular the construction of $\mathbf{Sym}^{(\ell)}$, have been used by Baumann and Hohlweg [2], whose paper provide detailed proofs. Hence, we shall only include the proofs which cannot be found in their paper. In particular, we propose an alternative approach to the internal product, which is introduced by a duality argument, and derive its main properties from those of the dual coproduct.

Acknowledgements. This project has been partially supported by the grant ANR-06-BLAN-0380. The authors would also like to thank the contributors of the MuPAD project, and especially of the combinat part, for providing the development environment for their research (see [15] for an introduction to MuPAD-Combinat).

2. BACKGROUND AND NOTATIONS

We first explain how to adapt the classical definitions and operations to the ℓ -colored case.

2.1. Colored alphabets. We shall start with an ℓ -colored alphabet

$$(1) \quad \mathbf{A} = A^0 \sqcup A^1 \sqcup \dots \sqcup A^{\ell-1},$$

such that all A^i are of the same cardinality N , which will be assumed to be infinite in the sequel. Let C be the alphabet $\{c_0, \dots, c_{\ell-1}\}$ and A be the auxiliary ordered alphabet $\{1, 2, \dots\}$ (the letter C stands for *colors* and A for *alphabet*) so that $\mathbf{A}$ can be identified with the cartesian product $A \times C$:

$$(2) \quad \mathbf{A} \simeq A \times C = \{(a, c), a \in A, c \in C\}.$$

A colored letter (i, c) will be denoted in bold type $\mathbf{i}$. Given two colored words, their *concatenation* is obtained by concatenating separately the elements coming from A and from C . We will sometimes allow $\ell = \infty$.

2.2. Colored standardization. Let w be a word in $\mathbf{A}$, represented as (v, u) with $v \in A^*$ and $u \in C^*$. Then the *colored standardized word* $\mathbf{Std}(w)$ of w is

$$(3) \quad \mathbf{Std}(w) := (\mathbf{Std}(v), u),$$

where $\mathbf{Std}(v)$ is the usual standardization on words.

Recall that the standardization process sends a word v of length n to a permutation $\mathbf{Std}(v) \in \mathfrak{S}_n$, called the *standardized* of v , defined as the permutation obtained by iteratively scanning v from left to right, and labelling $1, 2, \dots$ the occurrences of its smallest letter, then numbering the occurrences of the next one, and so on. Alternatively, $\mathbf{Std}(v)$ is the permutation having the same inversions as v .

For example, $\mathbf{Std}(abcadbdaa) = 157286934$:

$$(4) \quad \begin{array}{cccccccc} a & b & c & a & d & b & d & a & a \\ a_1 & b_5 & c_7 & a_2 & d_8 & b_6 & d_9 & a_3 & a_4 \\ 1 & 5 & 7 & 2 & 8 & 6 & 9 & 3 & 4 \end{array}$$

so that

$$(5) \quad \mathbf{Std}(abcadbdaa, 144120100) = (157286934, 144120100).$$

2.3. Colored shifted operations. For an element $v = (v_1, v_2, \dots)$ of A and an integer k , denote by $v[k]$ the *shifted word* $(v_1 + k) \cdots (v_n + k)$, e.g., $312[4] = 756$. Given a colored word $\alpha = (\alpha, u)$, we set $\alpha[k] = (\alpha[k], u)$.

The *shifted concatenation* of two words v and v' is defined by

$$(6) \quad v \bullet v' := v \cdot v'[k]$$

where k is the length of v .

The *shifted concatenation of two colored words* $(v, c), (v', c')$ is defined by

$$(7) \quad (v, c) \bullet (v', c') := (v \cdot v'[k], c \cdot c').$$

For example,

$$(8) \quad (13241, 00322) \bullet (12, 23) = (1324167, 0032223).$$

Finally, recall that the *shuffle product* of two words au and bv is defined by

$$(9) \quad au \sqcup bv = a(u \sqcup bv) + b(au \sqcup v),$$

where a and b are letters and u and v are words, with the initial conditions

$$(10) \quad u \sqcup \epsilon = \epsilon \sqcup u = u, \quad \epsilon \text{ being the empty word.}$$

This extends to the colored case, considering colored words as the concatenation of biletters.

The *shifted shuffle product* is

$$(11) \quad u \uplus v := u \sqcup v[k],$$

where k is the size of u .

3. FREE QUASI-SYMMETRIC FUNCTIONS OF LEVEL ℓ

3.1. $\mathbf{FQSym}^{(\ell)}$ and $\mathbf{FQSym}^{(\Gamma)}$. A *colored permutation* is a pair (σ, u) , with $\sigma \in \mathfrak{S}_n$ and $u \in C^n$, the integer n being the *size* of this permutation.

Definition 3.1. The dual free colored quasi-ribbon $\mathbf{G}_{\sigma,u}$ labelled by a colored permutation (σ, u) of size n is the noncommutative polynomial

$$(12) \quad \mathbf{G}_{\sigma,u} := \sum_{w \in A^n; \mathbf{Std}(w) = (\sigma,u)} w \in \mathbb{Z}\langle \mathbf{A} \rangle.$$

Recall that the *convolution* of two permutations σ and μ is the set $\sigma * \mu$ (identified with the formal sum of its elements) of permutations τ such that the standardized word of the $|\sigma|$ first letters of τ is σ and the standardized word of the remaining letters of τ is μ (see [39]). We then have:

Theorem 3.2. Let (σ', u') and (σ'', u'') be colored permutations. Then

$$(13) \quad \mathbf{G}_{\sigma',u'} \mathbf{G}_{\sigma'',u''} = \sum_{\sigma \in \sigma' * \sigma''} \mathbf{G}_{\sigma,u' \cdot u''}.$$

Therefore, the dual free colored quasi-ribbons span a $\mathbb{Z}$ -subalgebra $\mathbf{FQSym}^{(\ell)}$ of the free associative algebra.

Proof – This is immediate from the product of the usual free quasi-symmetric functions $\mathbf{G}_\sigma$:

$$(14) \quad \mathbf{G}_{\sigma'} \mathbf{G}_{\sigma''} = \sum_{\sigma \in \sigma' * \sigma''} \mathbf{G}_\sigma.$$

■

Note that all colored permutations indexing a product of $\mathbf{G}$ have given colors at the same places. For example,

$$(15) \quad \begin{aligned} \mathbf{G}_{(21,41)} \mathbf{G}_{(12,31)} = & \mathbf{G}_{(2134,4131)} + \mathbf{G}_{(3124,4131)} + \mathbf{G}_{(4123,4131)} \\ & + \mathbf{G}_{(3214,4131)} + \mathbf{G}_{(4213,4131)} + \mathbf{G}_{(4312,4131)}. \end{aligned}$$

One can define a coproduct by the usual trick of sums of alphabets: observe that we only need a total order on A to define the colored standardisation, so that taking two isomorphic copies A' and A'' of A , we define $\mathbf{A}' \oplus \mathbf{A}''$ as $(A' \oplus A'') \times C$, where

$A' \oplus A''$ denotes the ordered sum. Assuming furthermore that $\mathbf{A}'$ and $\mathbf{A}''$ commute, we identify $f(\mathbf{A}')g(\mathbf{A}'')$ with $f \otimes g$ and define a coproduct by:

$$(16) \quad \Delta f(\mathbf{A}) = f(\mathbf{A}' \oplus \mathbf{A}'').$$

By construction, this is an algebra morphism from $\mathbf{FQSym}^{(\ell)}$ to $\mathbf{FQSym}^{(\ell)} \otimes \mathbf{FQSym}^{(\ell)}$, so that

Theorem 3.3. $\mathbf{FQSym}^{(\ell)}$ is a graded connected bialgebra. Hence, it is a Hopf algebra. The coproduct is given by

$$(17) \quad \Delta \mathbf{G}_{\sigma, u} := \sum_{\substack{(\sigma', \sigma'', u', u'') \\ (\sigma, u) \in (\sigma', u') \uplus (\sigma'', u'')}} \mathbf{G}_{(\sigma', u')} \otimes \mathbf{G}_{(\sigma'', u'')}.$$

For example,

$$(18) \quad \begin{aligned} \Delta \mathbf{G}_{3142, 2412} &= 1 \otimes \mathbf{G}_{3142, 2412} + \mathbf{G}_{1,4} \otimes \mathbf{G}_{231, 212} + \mathbf{G}_{12, 42} \otimes \mathbf{G}_{12, 21} \\ &\quad + \mathbf{G}_{312, 242} \otimes \mathbf{G}_{1,1} + \mathbf{G}_{3142, 2412} \otimes 1. \end{aligned}$$

Proof – This is again immediate from the coproduct of the usual free quasi-symmetric functions $\mathbf{G}_\sigma$:

$$(19) \quad \Delta \mathbf{G}_\sigma = \sum_{\sigma \in \sigma' \uplus \sigma''} \mathbf{G}_{\sigma'} \otimes \mathbf{G}_{\sigma''}.$$

■

3.2. Duality in $\mathbf{FQSym}^{(\ell)}$.

Definition 3.4. The free ℓ -quasi-ribbon $\mathbf{F}_{\sigma, u}$ labelled by a colored permutation (σ, u) is the noncommutative polynomial

$$(20) \quad \mathbf{F}_{\sigma, u} := \mathbf{G}_{\sigma^{-1}, u \cdot \sigma^{-1}},$$

where the action of a permutation on the right of a word permutes the positions of the letters of the word.

For example,

$$(21) \quad \mathbf{F}_{3142, 2142} = \mathbf{G}_{2413, 1422}.$$

The product and coproduct of the $\mathbf{F}_{\sigma, u}$ can be easily described in terms of shifted shuffle and deconcatenation of colored permutations.

Theorem 3.5. Let σ' and σ'' be two colored permutations. Then

$$(22) \quad \mathbf{F}_{\sigma'} \mathbf{F}_{\sigma''} = \sum_{\sigma \in \sigma' \uplus \sigma''} \mathbf{F}_\sigma,$$

and

$$(23) \quad \Delta \mathbf{F}_\sigma = \sum_{\substack{w', w'' \\ \sigma = w' \cdot w''}} \mathbf{F}_{\text{Std}(w')} \otimes \mathbf{F}_{\text{Std}(w'')}.$$

Proof – Without colors, these formulas are the usual product of coproduct formulas of the $\mathbf{F}$ in $\mathbf{FQSym}$. With colors, one just has to observe that colors follow the letter to which they are attached. ■

Note that all colored permutations indexing a product of $\mathbf{F}$ have given colors associated with the same values, which is consistent with the corresponding remark on the $\mathbf{G}$ since places and values are exchanged when taking the inverse of a permutation.

For example, compare the following with Equation (15):

$$(24) \quad \begin{aligned} \mathbf{F}_{(21,14)} \mathbf{F}_{(12,31)} = & \mathbf{F}_{(2134,1431)} + \mathbf{F}_{(2314,1341)} + \mathbf{F}_{(2341,1314)} \\ & + \mathbf{F}_{(3214,3141)} + \mathbf{F}_{(3241,3114)} + \mathbf{F}_{(3421,3114)}. \end{aligned}$$

Here is an example of coproduct on the $\mathbf{F}$ basis:

$$(25) \quad \begin{aligned} \Delta \mathbf{F}_{(23514,14212)} = & 1 \otimes \mathbf{F}_{(23514,14212)} + \mathbf{F}_{(1,1)} \otimes \mathbf{F}_{(2413,4212)} + \mathbf{F}_{(12,14)} \otimes \mathbf{F}_{(312,212)} \\ & + \mathbf{F}_{(123,142)} \otimes \mathbf{F}_{(12,12)} + \mathbf{F}_{(2341,1421)} \otimes \mathbf{F}_{(1,12)} + \mathbf{F}_{(23514,14212)} \otimes 1. \end{aligned}$$

Let us define a scalar product on $\mathbf{FQSym}^{(\ell)}$ by

$$(26) \quad \langle \mathbf{F}_{\sigma,u}, \mathbf{G}_{\sigma',u'} \rangle := \delta_{\sigma,\sigma'} \delta_{u,u'},$$

where δ is the Kronecker symbol.

Theorem 3.6. *For any $U, V, W \in \mathbf{FQSym}^{(\ell)}$,*

$$(27) \quad \langle \Delta U, V \otimes W \rangle = \langle U, VW \rangle,$$

so that, $\mathbf{FQSym}^{(\ell)}$ is self-dual: the map $\mathbf{F}_{\sigma,u} \mapsto \mathbf{G}_{\sigma,u}^$ is an isomorphism from $\mathbf{FQSym}^{(\ell)}$ to its graded dual.*

Proof – Straightforward from Theorem 3.5. ■

Note 3.7. Let ϕ be any bijection from C to C , extended to words by concatenation. Then if one defines the free ℓ -quasi-ribbon as

$$(28) \quad \mathbf{F}_{\sigma,u} := \mathbf{G}_{\sigma^{-1}, \phi(u) \cdot \sigma^{-1}},$$

the previous theorems remain valid since one only permutes the labels of the basis $(\mathbf{F}_{\sigma,u})$. Moreover, if C has a semigroup structure, the colored permutations $(\sigma, u) \in \mathfrak{S}_n \times C^n$ can be interpreted as elements of the semi-direct product $H_n := \mathfrak{S}_n \ltimes C^n$ with multiplication rule

$$(29) \quad (\sigma; c_1, \dots, c_n) \cdot (\tau; d_1, \dots, d_n) := (\sigma\tau; c_{\tau(1)}d_1, \dots, c_{\tau(n)}d_n).$$

In furthermore C is a group, one can choose $\phi(\gamma) := \gamma^{-1}$ and define the scalar product as before, so that the adjoint basis of the $(\mathbf{G}_h)$ becomes $\mathbf{F}_h := \mathbf{G}_{h^{-1}}$. In the sequel, we will be mainly interested in the cases $C := \mathbb{Z}/\ell\mathbb{Z}$, and we will indeed make that choice for ϕ whenever C is a group.

3.3. Algebraic structure of $\mathbf{FQSym}^{(\ell)}$. Recall that a permutation σ of size n is *connected* [27, 4] if, for any $i < n$, the set $\{\sigma_1, \dots, \sigma_i\}$ is different from $\{1, \dots, i\}$.

We denote by $\mathcal{C}$ the set of connected permutations, and by $c_n := |\mathcal{C}_n|$ the number of such permutations in $\mathfrak{S}_n$. For later reference, we recall that the generating series of c_n is Sequence A003319 of [41]:

$$(30) \quad c(t) := \sum_{n \geq 1} c_n t^n = 1 - \left(\sum_{n \geq 0} n! t^n \right)^{-1} = t + t^2 + 3t^3 + 13t^4 + 71t^5 + 461t^6 + O(t^7).$$

Let the *connected colored permutations* be the (σ, u) with σ connected and u arbitrary. Their generating series is given by $c(\ell t)$.

From [4], we immediately get

Proposition 3.8. $\mathbf{FQSym}^{(\ell)}$ is free over the set $\mathbf{F}_{\sigma,u}$ (or $\mathbf{G}_{\sigma,u}$), where (σ, u) is connected. ■

For example, the generating series of the algebraic generators of $\mathbf{FQSym}^{(2)}$ is

$$(31) \quad 2t + 4t^2 + 24t^3 + 208t^4 + 2272t^5 + 29504t^6 + 441216t^7 + \dots$$

3.4. Primitive elements of $\mathbf{FQSym}^{(\ell)}$. Let $\mathcal{L}^{(\ell)}$ be the primitive Lie algebra of $\mathbf{FQSym}^{(\ell)}$. Since Δ is not cocommutative, $\mathbf{FQSym}^{(\ell)}$ cannot be the universal enveloping algebra of $\mathcal{L}^{(\ell)}$. But since it is cofree, it is, according to [23], the universal enveloping dipterous algebra of its primitive part $\mathcal{L}^{(\ell)}$.

Let $\mathbf{G}^{\sigma,u}$ be the multiplicative basis defined by $\mathbf{G}^{\sigma,u} = \mathbf{G}_{\sigma_1, u_1} \cdots \mathbf{G}_{\sigma_r, u_r}$ where $(\sigma, u) = (\sigma_1, u_1) \bullet \cdots \bullet (\sigma_r, u_r)$ is the unique maximal factorization of $(\sigma, u) \in \mathfrak{S}_n \times C^n$ into connected colored permutations.

Proposition 3.9. Let $\mathbf{V}_{\sigma,u}$ be the adjoint basis of $\mathbf{G}^{\sigma,u}$. Then, the family $(\mathbf{V}_{\alpha,u})_{\alpha \in \mathcal{C}}$ is a basis of $\mathcal{L}^{(\ell)}$. In particular, we have $\dim \mathcal{L}_n^{(\ell)} = \ell^n c_n$. Moreover, $\mathcal{L}^{(\ell)}$ is free.

Proof – The first part of the statement follows from [4]. The second part comes from the fact that $\mathbf{FQSym}^{(\ell)}$ is bidendriform (Theorem 3.10 below). ■

For example, since $\mathcal{L}^{(\ell)}$ is free, the generating series by degree of its generators is (with $\ell = 2$):

$$(32) \quad \begin{aligned} 1 - \prod_{n \geq 1} (1 - t^n)^{\ell^n c_n} &= 1 - (1 - t)^2 (1 - t^2)^4 (1 - t^3)^{24} \dots \\ &= 2t + 3t^2 + 16t^3 + 158t^4 + 1796t^5 \\ &\quad + 24250t^6 + 372656t^7 + \dots \end{aligned}$$

and the Hilbert series of the universal enveloping algebra of $\mathbf{FQSym}^{(\ell)}$ (its domain of cocommutativity) is, again with $\ell = 2$,

$$(33) \quad \begin{aligned} \prod_{n \geq 1} (1 - t^n)^{-\ell^n c_n} &= 1 + 2t + 7t^2 + 36t^3 + 283t^4 + 2898t^5 + 36169t^6 \\ &\quad + 524976t^7 + \dots \end{aligned}$$

3.5. Dendriform structure of $\mathbf{FQSym}^{(\ell)}$. Foissy introduced the notion of *bidendriform* bialgebras [6], generalizing the notion of *dendriform* algebras (cf. [24]) and proved some conjectures about $\mathbf{FQSym}$, presented in [4]. We shall adapt this technology to the colored case. We shall not recall all the theory, since complete details can be found in [6].

Recall that the generators of $\mathbf{FQSym}^{(\ell)}$ as a dendriform algebra are called *totally primitive elements* [6] and that their generating series is given by

$$(34) \quad \text{TP} := \frac{\text{PQ} - 1}{\text{PQ}^2},$$

where PQ is the generating series of $\mathbf{FQSym}^{(\ell)}$.

Theorem 3.10. *The algebra $\mathbf{FQSym}^{(\ell)}$ has a structure of bidendriform bialgebra [6], hence is free as a Hopf algebra and as a dendriform algebra, cofree, self-dual, and its primitive Lie algebra is free.*

Moreover, the totally primitive elements of $\mathbf{FQSym}^{(\ell)}$ are the totally primitive elements of $\mathbf{FQSym}$ with any coloring.

Proof – It has been done by Foissy in [6] in the case of $\mathbf{FQSym}$. But since the dendriform and codendriform structure do not involve the color alphabet C , the property is true in this case as well.

Similarly, colors do not play any role in determining if a given element is (totally) primitive or not. ■

For example, the dendriform generators of $\mathbf{FQSym}$ have as degree generating series

$$(35) \quad \sum_i \text{dg}_i t^i = t + t^3 + 6t^4 + 39t^5 + 284t^6 + 2305t^7 + \dots$$

so that the dendriform generators of $\mathbf{FQSym}^{(2)}$ have as degree generating series 2^idg_i :

$$(36) \quad 2t + 8t^3 + 96t^4 + 1248t^5 + 18176t^6 + 295040t^7 + \dots,$$

Note that there cannot be any relation, even dendriform relations, among the elements $\mathbf{F}_{1,c}$ where $c \in C$, so that $\mathbf{FQSym}^{(\ell)}$ contains the free dendriform algebra $\mathbf{PBT}^{(\ell)}$ on ℓ generators.

3.6. Internal product of $\mathbf{FQSym}^{(\ell)}$. When C is a semigroup, an internal product can be defined on $\mathbf{FQSym}^{(\ell)}$ by

$$(37) \quad \mathbf{F}_{\sigma,u} * \mathbf{F}_{\tau,v} = \mathbf{F}_{\mu,w}$$

where (μ, w) is the product $(\sigma, u) \cdot (\tau, v)$ in the wreath product, defined by formula (29), that is

$$(38) \quad \mathbf{F}_{(\sigma,u)} * \mathbf{F}_{(\tau,v)} = \mathbf{F}_{(\sigma\tau, (u\tau) \cdot v)},$$

where $u\tau$ is the word $u_{\tau_1} \dots u_{\tau_n}$ and $u \cdot v$ is the componentwise product defined by $(u_1 v_1, \dots, u_n v_n)$.

For example, if the color group is $\mathbb{Z}$

$$(39) \quad \mathbf{F}_{(1324,1011)} * \mathbf{F}_{(2413,3200)} = \mathbf{F}_{(3412,3311)}.$$

$$(40) \quad \mathbf{F}_{(165324,102011)} * \mathbf{F}_{(625413,322011)} = \mathbf{F}_{(462315,423023)}.$$

This can be reduced to any $\mathbb{Z}/\ell\mathbb{Z}$, *e.g.*, with $\ell = 3$,

$$(41) \quad \mathbf{F}_{(165324,102011)} * \mathbf{F}_{(625413,022011)} = \mathbf{F}_{(462315,120020)}.$$

In the $\mathbf{G}$ basis, one has

$$(42) \quad \mathbf{G}_{(\sigma,u)} * \mathbf{G}_{(\tau,v)} = \mathbf{G}_{(\tau\sigma,u.(v\sigma))},$$

4. NONCOMMUTATIVE SYMMETRIC FUNCTIONS OF LEVEL ℓ

4.1. ℓ -partite numbers. Following McMahon [26], we define an ℓ -partite number $\mathbf{n} = (n_1, \dots, n_\ell)$ as a column vector in $\mathbb{N}^\ell$, and a *vector composition of $\mathbf{n}$ of weight $|\mathbf{n}| := \sum_1^\ell n_i$ and length m* as a $\ell \times m$ matrix $\mathbf{I}$ of nonnegative integers, with row sums vector $\mathbf{n}$ and no zero column.

For example,

$$(43) \quad \mathbf{I} = \begin{pmatrix} 1 & 0 & 2 & 1 \\ 0 & 3 & 1 & 1 \\ 4 & 2 & 1 & 3 \end{pmatrix}$$

is a vector composition (or a 3-composition, for short) of the 3-partite number $\begin{pmatrix} 4 \\ 5 \\ 10 \end{pmatrix}$ of weight 19 and length 4.

For each $\mathbf{n} \in \mathbb{N}^\ell$ of weight $|\mathbf{n}| = n$, we define a *level ℓ complete homogeneous noncommutative symmetric function* as

$$(44) \quad S_{\mathbf{n}} := \sum_{u; |u|_i = n_i} \mathbf{G}_{1\dots n, u}.$$

It is the sum of all possible colorings of the identity permutation with n_i occurrences of color i for each i .

4.2. The Hopf algebra $\mathbf{Sym}^{(\ell)}$. Let $\mathbf{Sym}^{(\ell)}$ be the subalgebra of $\mathbf{FQSym}^{(\ell)}$ generated by the $S_{\mathbf{n}}$ (with the convention $S_0 = 1$). The Hilbert series of $\mathbf{Sym}^{(\ell)}$ is easily found to be

$$(45) \quad S_\ell(t) := \sum_n \dim \mathbf{Sym}_n^{(\ell)} t^n = \frac{(1-t)^\ell}{2(1-t)^\ell - 1}.$$

For example, with $\ell = 2$, one has

$$(46) \quad S_2(t) := 1 + 2t + 7t^2 + 24t^3 + 82t^4 + 280t^5 + 956t^6 + 3264t^7 + \dots$$

which is Sequence A003480 of [41].

For general ℓ , it is well-known in the combinatorial folklore (and easy to prove by means of generating series expansions) that, for all $n \geq 1$,

$$(47) \quad \text{ncs}_n(\ell) := n! \dim(\mathbf{Sym}_n^{(\ell)}) = \sum_{k=1}^n S(n, k) p_k \ell^k$$

where $S(n, k)$ is the sequence of absolute values of Stirling numbers of the first kind (sequence A130534 of [41]) and p_k is the sequence of ordered Bell numbers (also known as packed words or preferential arrangements, Sequence A000670 of [41]).

Theorem 4.1. $\mathbf{Sym}^{(\ell)}$ is free over the set $\{S_{\mathbf{n}}, |\mathbf{n}| > 0\}$, so that a linear basis is given by

$$(48) \quad S^{\mathbf{I}} = S_{\mathbf{i}_1} \cdots S_{\mathbf{i}_m},$$

where $\mathbf{i}_1, \dots, \mathbf{i}_m$ are the columns of $\mathbf{I}$.

Moreover, $\mathbf{Sym}^{(\ell)}$ is a Hopf subalgebra of $\mathbf{FQSym}^{(\ell)}$ and the coproduct of the generators is given by

$$(49) \quad \Delta S_{\mathbf{n}} = \sum_{\mathbf{i}+\mathbf{j}=\mathbf{n}} S_{\mathbf{i}} \otimes S_{\mathbf{j}},$$

where the sum $\mathbf{i} + \mathbf{j}$ is taken in the space $\mathbb{N}^l$. In particular, $\mathbf{Sym}^{(\ell)}$ is cocommutative.

Proof – Consider a linear relation between the $S^{\mathbf{I}}$. Since without colors the algebra is free (it is $\mathbf{Sym}$), the linear relation splits into many linear relations involving terms of the form $S^{\mathbf{i}_1 \dots \mathbf{i}_r}$ with $|\mathbf{i}_1|, \dots, |\mathbf{i}_r|$ fixed. But there are no relations of this form, thanks to the definition of the $\mathbf{G}_{\sigma}$ as a sum of colored words.

Given the coproduct of the $\mathbf{G}_{\sigma}$ of $\mathbf{FQSym}^{(\ell)}$ and since the $S_{\mathbf{n}}$ are sums of $\mathbf{G}$, the coproduct of an $S_{\mathbf{n}}$ amounts to unshuffling the color words, whence their coproduct formula. ■

For example,

$$(50) \quad \begin{aligned} \Delta S \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} = & S \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} \otimes S \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} + S \begin{pmatrix} 0 \\ 0 \\ 2 \end{pmatrix} \otimes S \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + S \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \otimes S \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \\ & + S \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \otimes S \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + S \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \otimes S \begin{pmatrix} 0 \\ 0 \\ 2 \end{pmatrix} + S \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \otimes S \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}. \end{aligned}$$

4.3. Algebraic structure of $\mathbf{Sym}^{(\ell)}$. The number of generators of $\mathbf{Sym}^{(\ell)}$ of degree n is given by the number of ℓ -partite numbers of total sum n . So its generating series is

$$(51) \quad (1 - t)^{-\ell} - 1 = \sum_{n \geq 1} \binom{\ell + n - 1}{n} t^n.$$

We shall denote by $G(\ell)$ the set of nonzero ℓ -partite numbers.

4.4. Primitive elements of $\mathbf{Sym}^{(\ell)}$. $\mathbf{Sym}^{(\ell)}$ being a graded connected cocommutative Hopf algebra, it follows from the Cartier-Milnor-Moore theorem that it is the universal enveloping algebra of $L^{(\ell)}$:

$$(52) \quad \mathbf{Sym}^{(\ell)} = U(L^{(\ell)}),$$

where $L^{(\ell)}$ is the Lie algebra of its primitive elements. Let us now prove

Theorem 4.2. *As a graded Lie algebra, the primitive Lie algebra $L^{(\ell)}$ of $\mathbf{Sym}^{(\ell)}$ is free over a set indexed by $G(\ell)$.*

Proof – If $L^{(\ell)}$ is free, by standard arguments on generating series, the number of generators of $L^{(\ell)}$ in degree n must be the number of algebraic generators of $\mathbf{Sym}^{(\ell)}$ in degree n , parametrized for example by $G(\ell)$. We will now show that $L^{(\ell)}$ has at least this number of generators and that those generators are algebraically independent, determining completely the dimensions of the homogeneous components $L_n^{(\ell)}$ of $L^{(\ell)}$ whose generating series begins by

$$(53) \quad t + \binom{\ell+1}{2} t^2 + \left(\binom{\ell+1}{2} + \binom{\ell+2}{3} \right) t^3 + \dots$$

Following Reutenauer [39] p. 58, denote by π_1 the Eulerian idempotent, that is, the endomorphism of $\mathbf{Sym}^{(\ell)}$ defined by $\pi_1 = \log^*(Id)$. It is obvious, thanks to the definition of $S_{\mathbf{p}}$ that

$$(54) \quad \pi_1(S_{\mathbf{p}}) = S_{\mathbf{p}} + \dots,$$

where the dots stand for terms $S_{\mathbf{I}}$ such that $\mathbf{I}$ are vector compositions with strictly more than one column. Since the $S_{\mathbf{p}}$ where $\mathbf{p}$ is a ℓ -partite number are algebraically independent, the dimension of $L_n^{(\ell)}$ is at least equal to the cardinality of the elements of $G(\ell)$ of size n . So $L^{(\ell)}$ is indeed free over a set of primitive elements parametrized by $G(\ell)$. ■

For example, with $\ell = 2$, the generating series of the dimensions of $L^{(\ell)}$ is

$$(55) \quad 1 + 2t + 4t^2 + 12t^3 + 31t^4 + 92t^5 + 256t^6 + 772t^7 + \dots$$

With $\ell = 3$, one finds

$$(56) \quad 1 + 3t + 9t^2 + 36t^3 + 132t^4 + 534t^5 + 2140t^6 + 8982t^7 + \dots$$

4.5. Duality in $\mathbf{Sym}^{(\ell)}$. Recall that the underlying colored alphabet $\mathbf{A}$ can be seen as $A^0 \sqcup \dots \sqcup A^{\ell-1}$, with $A^i = \{a_j^{(i)}, j \geq 1\}$. Let $\mathbf{x} = (x^{(0)}, \dots, x^{(\ell-1)})$, where the $x^{(i)}$ are ℓ commuting variables. In terms of $\mathbf{A}$, the generating function of the complete functions can be written as

$$(57) \quad \sigma_{\mathbf{x}}(\mathbf{A}) = \prod_{i \geq 0}^{\rightarrow} \left(1 - \sum_{0 \leq j \leq \ell-1} x^{(j)} a_i^{(j)} \right)^{-1} = \sum_{\mathbf{n}} S_{\mathbf{n}}(\mathbf{A}) \mathbf{x}^{\mathbf{n}},$$

where $\mathbf{x}^{\mathbf{n}} = (x^{(0)})^{n_0} \dots (x^{(\ell-1)})^{n_{\ell-1}}$.

This realization gives rise to a Cauchy formula (see [20] for the $l = 1$ case), which in turn allows one to identify the dual of $\mathbf{Sym}^{(\ell)}$ with an algebra introduced by S. Poirier in [35]. It is detailed in the following section.

Note that $\mathbf{Sym}^{(\ell)}$ is the natural noncommutative analog of McMahon's algebra of multisymmetric functions [26, 10].

5. QUASI-SYMMETRIC FUNCTIONS OF LEVEL ℓ

5.1. Cauchy formula of level ℓ . Let $\mathbb{X} = X^0 \sqcup \cdots \sqcup X^{\ell-1}$, where $X^i = \{x_j^{(i)}, j \geq 1\}$, be an ℓ -colored alphabet of commutative variables, also commuting with $\mathbf{A}$. Imitating the level 1 case (see [4]), we define the Cauchy kernel

$$(58) \quad K(\mathbb{X}, \mathbf{A}) = \prod_{j \geq 1}^{\rightarrow} \sigma_{(x_j^{(0)}, \dots, x_j^{(\ell-1)})}(\mathbf{A}).$$

Expanding on the basis $S^{\mathbf{I}}$ of $\mathbf{Sym}^{(\ell)}$, we get as coefficients what can be called the *level ℓ monomial quasi-symmetric functions* $M_{\mathbf{I}}(\mathbb{X})$

$$(59) \quad K(\mathbb{X}, \mathbf{A}) = \sum_{\mathbf{I}} M_{\mathbf{I}}(\mathbb{X}) S^{\mathbf{I}}(\mathbf{A}),$$

defined by

$$(60) \quad M_{\mathbf{I}}(\mathbb{X}) = \sum_{j_1 < \cdots < j_m} \mathbf{x}_{j_1}^{\mathbf{i}_1} \cdots \mathbf{x}_{j_m}^{\mathbf{i}_m},$$

with $\mathbf{I} = (\mathbf{i}_1, \dots, \mathbf{i}_m)$.

These functions form a basis of a subalgebra $QSym^{(\ell)}$ of $\mathbb{K}[\mathbb{X}]$, which we shall call the *algebra of quasi-symmetric functions of level ℓ* .

5.2. Poirier's Quasi-symmetric functions. The functions $M_{\mathbf{I}}(\mathbb{X})$ can be recognized as a basis of one of the algebras introduced by Poirier: the $M_{\mathbf{I}}$ coincide with the $M_{(C,v)}$ defined in [35], p. 324, formula (1), up to indexation.

Following Poirier, we introduce the level ℓ quasi-ribbon functions by summing over an order on ℓ -compositions: an ℓ -composition C is finer than C' , and we write $C \leq C'$, if C' can be obtained by repeatedly summing up two consecutive columns of C such that no nonzero element of the left one is strictly below a nonzero element of the right one.

This order can be described in a much easier and natural way if one recodes an ℓ -composition $\mathbf{I}$ as a pair of words, the first one $d(\mathbf{I})$ being the set of sums of the elements of the first k columns of $\mathbf{I}$, the second one $c(\mathbf{I})$ being obtained by concatenating the words $i^{\mathbf{I}_{i,j}}$ while reading of $\mathbf{I}$ by columns, from top to bottom and from left to right. For example, the 3-composition of Equation (43) satisfies

$$(61) \quad d(\mathbf{I}) = \{5, 10, 14, 19\} \quad \text{and} \quad c(\mathbf{I}) = 13333 \, 22233 \, 1123 \, 12333.$$

Moreover, this recoding is a bijection if the two words $d(\mathbf{I})$ and $c(\mathbf{I})$ are such that the descent set of $c(\mathbf{I})$ is a subset of $d(\mathbf{I})$. The order previously defined on ℓ -compositions is in this context the inclusion order on sets d : $(d', c) \leq (d, c)$ iff $d' \subseteq d$.

It allows us to define the *level ℓ quasi-ribbon functions* $F_{\mathbf{I}}$ by

$$(62) \quad F_{\mathbf{I}} = \sum_{\mathbf{I}' \leq \mathbf{I}} M_{\mathbf{I}'}.$$

Notice that this last description of the order $\leq$ is reminiscent of the order $\leq'$ on descent sets used in the context of quasi-symmetric functions and non-commutative symmetric functions: more precisely, since it does not modify the word $c(\mathbf{I})$, the order $\leq$ restricted to ℓ -compositions of weight n amounts for ℓ^n copies of the order $\leq'$. The computation of its Möbius function is therefore straightforward.

Moreover, one can obtain the $F_{\mathbf{I}}$ as the commutative image of certain $\mathbf{F}_{\sigma, u}$: any pair (σ, u) such that σ has descent set $d(\mathbf{I})$ and $u = c(\mathbf{I})$ will do.

5.3. Coproducts and alphabets.

5.3.1. Recall that to define the $\mathbf{G}_{\sigma, u}(\mathbf{A})$ of an ℓ -colored alphabet $\mathbf{A} = A \times C$, we only need a total order on A . Hence, if B is another copy of A commuting with A , we can define $\mathbf{A} + \mathbf{B}$ as $(A + B) \times C$ where $A + B$ is the ordinal sum, and thus make sense of $\mathbf{G}_{\sigma, u}(\mathbf{A} + \mathbf{B})$.

As usual, we identify $F(\mathbf{A})G(\mathbf{B})$ with $F \otimes G$.

Lemma 5.1. *For any $F \in \mathbf{FQSym}^{(\ell)}$, $\mathbf{F}(\mathbf{A} + \mathbf{B}) \in \mathbf{FQSym}^{(\ell)} \otimes \mathbf{FQSym}^{(\ell)}$, and*

$$(63) \quad \mathbf{F}(\mathbf{A} + \mathbf{B}) = \Delta F,$$

where Δ is the coproduct defined by (17).

Proof – It is sufficient to show this for $\ell = 1$, which is done in [4]. ■

Let us observe that on this picture, it is clear that the restriction of Δ to $\mathbf{Sym}^{(\ell)}$ is dual to the product of $QSym^{(\ell)}$. By definition of the Cauchy kernel (58), we have

$$(64) \quad K(\mathbb{X}, \mathbf{A} + \mathbf{B}) = K(\mathbb{X}, \mathbf{A})K(\mathbb{X}, \mathbf{B}),$$

and by (59), this implies that Δ is dual of the multiplication of $QSym^{(\ell)}$.

5.3.2. The same description can be applied to the quasi-symmetric side. Let $\mathbb{X} = X \times C$ and $\mathbb{Y} = Y \times C$, where Y is a copy of X . Again, define $X + Y$ as the ordinal sum of X and Y , and $\mathbb{X} + \mathbb{Y} = (X + Y) \times C$.

Lemma 5.2. *The map $\nabla : F \mapsto F(\mathbb{X} + \mathbb{Y})$ defines a coproduct on $QSym^{(\ell)}$, which is dual to the product of $\mathbf{Sym}^{(\ell)}$.*

Proof – This follows from the identity

$$(65) \quad K(\mathbb{X} + \mathbb{Y}, \mathbf{A}) = K(\mathbb{X}, \mathbf{A})K(\mathbb{Y}, \mathbf{A}).$$

■

5.3.3. *The internal coproduct.* From now on, we assume that the color set C is an additive semigroup, such that every element $\gamma \in C$ has a finite number of decompositions $\gamma = \alpha + \beta$.

We define the C -product $\mathbb{T} = \mathbb{X} \times_C \mathbb{Y}$ of two C -colored alphabets by

$$(66) \quad \mathbb{T}^{(\gamma)} = \{t_{rs}^{(\gamma)} = \sum_{\alpha+\beta=\gamma} x_r^{(\alpha)} y_s^{(\beta)}\},$$

with the pairs (r, s) ordered lexicographically.

Proposition 5.3. *The map $\delta : F \mapsto F(\mathbb{X} \times_C \mathbb{Y})$ defines a coassociative coproduct on $QSym^{(\ell)}$.*

We define the internal product $*$ of $\mathbf{Sym}^{(\ell)}$ as the dual product of the map δ .

Proof – The coassociativity condition

$$(67) \quad (\delta \otimes Id) \circ \delta = (Id \otimes \delta) \circ \delta$$

translates as the associativity of the C -product

$$(68) \quad (\mathbb{X} \times_C \mathbb{Y}) \times_C \mathbb{Z} = \mathbb{X} \times_C (\mathbb{Y} \times_C \mathbb{Z})$$

which is clear since both sides are by definition

$$(69) \quad \left\{ t_{pqr}^\mu = \sum_{\alpha+\beta+\gamma=\mu} x_p^{(\alpha)} y_q^{(\beta)} z_r^{(\gamma)} \right\}$$

with the lexicographic order on triples (p, q, r) . ■

Example 5.4. With $\ell = 2$ and $C = \mathbb{Z}/2\mathbb{Z}$,

$$\begin{aligned} s \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} *_s s \begin{pmatrix} 0 & 2 \\ 1 & 0 \end{pmatrix} &= \mu_2 \left[\left(s \begin{pmatrix} 1 \\ 1 \end{pmatrix} \otimes s \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right) *_2 \Delta s \begin{pmatrix} 0 & 2 \\ 1 & 0 \end{pmatrix} \right] \\ &= \left(s \begin{pmatrix} 1 \\ 1 \end{pmatrix} *_s s \begin{pmatrix} 2 \\ 0 \end{pmatrix} \right) \left(s \begin{pmatrix} 0 \\ 1 \end{pmatrix} *_s s \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right) + \left(s \begin{pmatrix} 1 \\ 1 \end{pmatrix} *_s s \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right) \left(s \begin{pmatrix} 0 \\ 1 \end{pmatrix} *_s s \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right) \\ &= s \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} + s \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + s \begin{pmatrix} 0 & 0 & 0 \\ 1 & 1 & 1 \end{pmatrix}. \end{aligned}$$

5.3.4. *The splitting formula.* The definition of $*$ by duality with the C -product implies that it satisfies the splitting formula. We just need to check a trivial property:

Lemma 5.5. *The C -product is distributive over the colored ordinal sum:*

$$(70) \quad (\mathbb{X}' + \mathbb{X}'') \times_C \mathbb{Y} = \mathbb{X}' \times_C \mathbb{Y} + \mathbb{X}'' \times_C \mathbb{Y}.$$
■

Proposition 5.6. *Let $\mu_r : (\mathbf{Sym}^{(\ell)})^{\otimes r} \rightarrow \mathbf{Sym}^{(\ell)}$ be the product map. Let $\Delta^{(r)} : (\mathbf{Sym}^{(\ell)}) \rightarrow (\mathbf{Sym}^{(\ell)})^{\otimes r}$ be the r -fold coproduct, and $*_r$ be the extension of the internal product to $(\mathbf{Sym}^{(\ell)})^{\otimes r}$. Then, for $F_1, \dots, F_r$, and $G \in \mathbf{Sym}^{(\ell)}$,*

$$(71) \quad (F_1 \cdots F_r) * G = \mu_r[(F_1 \otimes \cdots \otimes F_r) *_r \Delta^{(r)} G].$$

Proof – It is enough to consider the case $r = 2$. Let $F \in QSym^{(\ell)}$ and $U, W, W \in \mathbf{Sym}^{(\ell)}$. We have, writing $\mathbb{X}\mathbb{Y} = \mathbb{X} \times_C \mathbb{Y}$ for short, and assuming duality between $\mathbb{X}$ and $\mathbf{A}$, $\mathbb{Y}$ and $\mathbf{B}$ and so on,

$$\begin{aligned} \langle F, (UV) * W \rangle &= \langle F(\mathbb{X}'\mathbb{X}''), (UV)(\mathbf{A}')W(\mathbf{A}'') \rangle \\ &= \langle F(\mathbb{X}' + \mathbb{Y}')\mathbb{X}'' \rangle, U(\mathbf{A}')V(\mathbf{B}')W(\mathbf{A}'') \rangle \\ &= \langle F(\mathbb{X}'\mathbb{X}'' + \mathbb{Y}'\mathbb{Y}''), U(\mathbf{A}')V(\mathbf{B}')W(\mathbf{A}'' + \mathbf{B}'') \rangle \\ &= \langle F(\mathbb{X}' + \mathbb{Y}'), U(\mathbf{A}')V(\mathbf{B}')W(\mathbf{A}' + \mathbf{B}') \rangle \\ &= \langle \nabla F, (U \otimes V) *_2 \Delta W \rangle = \langle F, \mu[(U \otimes V) *_2 \Delta W] \rangle. \end{aligned}$$

■

5.3.5. *Evaluation of internal products.* Let us start with the simplest case, $S_i * S_j$. The coefficient of $S^{\mathbf{K}}$ in this product is the coefficient of $M_i \otimes M_j$ in $\delta M_{\mathbf{K}}$, which is also the coefficient of $\mathbf{x}^i \mathbf{y}^j$ in $M_{\mathbf{K}}(\mathbf{x} \times_C \mathbf{y})$. This is zero if $\mathbf{K}$ has more than one column, so that

$$(72) \quad S_i * S_j = \sum_{\mathbf{n}} d_{ij}^{\mathbf{n}} S_{\mathbf{n}}$$

contains only one-part vector compositions.

Lemma 5.7. *If the color semigroup is $C = \mathbb{N}$, then the coefficient $d_{ij}^{\mathbf{n}}$ of $S_{\mathbf{n}}$ in $S_i * S_j$ is equal to the coefficient of the monomial symmetric function m_{μ} in the product $m_{\alpha} m_{\beta}$, where μ is the partition $(0^{n_0} 1^{n_1} \dots)$, $\alpha = (0^{i_0} 1^{i_1} \dots)$, $\beta = (0^{j_0} 1^{j_1} \dots)$, the monomial functions being taken over an alphabet of n letters, where $n = |\mathbf{i}|$.*

Proof – From (72), we only need to compute the coproducts $\delta M_{\mathbf{n}}$. For this, it is sufficient to use alphabets of the form $\mathbf{x} = \{x\} \otimes C$, $\mathbf{y} = \{y\} \otimes C$. Then,

$$(73) \quad M_{\mathbf{n}}(\mathbf{x} \times_C \mathbf{y}) = \prod_{k \geq 0} \left(\sum_{i+j=k} x^{(i)} y^{(j)} \right)^{n_k},$$

and we see that the coefficient of $M_i(\mathbf{x}) M_j(\mathbf{y}) = \mathbf{x}^i \mathbf{y}^j$ is the same as the coefficient of $h_{\alpha} \otimes h_{\beta}$ in

$$(74) \quad \prod_{k \geq 0} \left(\sum_{i+j=k} h_i \otimes h_j \right)^{n_k} = \Delta(h_0^{n_0} h_1^{n_1} \dots) = \Delta h_{\mu}.$$

■

For example,

$$(75) \quad S^{\binom{2}{2}} * S^{\binom{3}{1}} = 3 S^{\binom{1}{3}} + S^{\binom{2}{1}}.$$

This result is compatible with the fact that

$$(76) \quad m_{11}m_1 = 3m_{111} + m_{21}.$$

As another example,

$$(77) \quad S^{\binom{0}{2}} * S^{\binom{1}{1}} = S^{\binom{0}{1}} + 2 S^{\binom{0}{2}} + 2 S^{\binom{0}{1}}.$$

One can then check that the previous result amounts to selecting the partitions of size at most 3 in

$$(78) \quad m_{211}m_{21} = m_{421} + 2m_{331} + 2m_{322} + \dots$$

Together with the splitting formula (71), this result determines all the products $S^{\mathbf{I}} * S^{\mathbf{J}}$, since one also has

$$(79) \quad S^{\mathbf{i}} * S^{\mathbf{J}} = S^{\mathbf{J}} * S^{\mathbf{i}}$$

thanks to the isomorphism of ordered colored alphabets

$$(80) \quad \mathbf{x} \times_C \mathbf{Y} \simeq \mathbf{Y} \times_C \mathbf{x},$$

where $\mathbf{x} = \{x\} \otimes C$.

When the color group is $\mathbb{Z}/\ell\mathbb{Z}$, the result is obtained by reduction modulo ℓ , *e.g.*, with $\ell = 3$, we get from example (77)

$$(81) \quad S^{\binom{0}{2}} * S^{\binom{1}{1}} = 2 S^{\binom{0}{1}} + 2 S^{\binom{2}{1}} + 2 S^{\binom{1}{2}}.$$

Note that the coefficient of an $S^{\mathbf{I}}$ can change when computing modulo ℓ : for all pairs of parts k and k' added together, a factor $\binom{k+k'}{k}$ appears.

5.4. Generalized descent algebras. In a preliminary draft of this work [32], we introduced the internal product in a different way. Assuming that C has a semigroup structure, we regard colored permutations as elements of the wreath product $H = C \wr \mathfrak{S}_n$, and for $h', h'' \in H$, we set

$$(82) \quad \mathbf{G}_{h'} *' \mathbf{G}_{h''} = \mathbf{G}_{h''h'}$$

(opposite law, as in the classical case of **Sym**).

Theorem 5.8. *Let C be a commutative semigroup.*

- (1) $\mathbf{Sym}_n^{(C)}$ is a subalgebra of $\mathbf{FQSym}^{(C)}$, for the operation $*'$ defined previously.
- (2) The restriction of $*'$ to $\mathbf{Sym}_n^{(C)}$ satisfies the splitting formula (71).
- (3) $S^{\mathbf{i}} *' S^{\mathbf{j}}$ is given by Lemma 5.7.
- (4) $S^{\mathbf{i}} *' S^{\mathbf{J}} = S^{\mathbf{J}} *' S^{\mathbf{i}}$.

Proof – The proofs of (1) and (2) can be found in [2]. (3) and (4) follow from the definition in (44) and the internal product on $\mathbf{FQSym}^{(C)}$ given by (38). ■

This provides an analogue of Solomon's descent algebra for the wreath product $C \wr \mathfrak{S}_n$. Note that the definition remains valid for $C = \mathbb{Z}$, so that we get a descent algebra for the (extended) affine Weyl groups of type A , $\hat{\mathfrak{S}}_n = \mathbb{Z} \wr \mathfrak{S}_n$.

5.5. Ordinary multi-symmetric functions. A consequence of the above results is that the algebra $Sym^{(\ell)}$ of ordinary (commutative) multi-symmetric functions admits an internal product. If we denote by $\underline{F} \in Sym^{(\ell)}$ the commutative image of $F \in \mathbf{Sym}^{(\ell)}$, we have

$$(83) \quad \underline{F} * \underline{G} = \underline{F'} * \underline{G'} \quad \text{as soon as } \underline{F} = \underline{F'} \text{ and } \underline{G} = \underline{G'}.$$

For example,

$$(84) \quad \begin{aligned} S \begin{pmatrix} 0 & 3 \\ 1 & 1 \end{pmatrix} * S \begin{pmatrix} 1 & 2 \\ 1 & 1 \end{pmatrix} = & S \begin{pmatrix} 0 & 0 & 2 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} + S \begin{pmatrix} 0 & 0 & 2 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{pmatrix} + S \begin{pmatrix} 0 & 1 & 2 \\ 0 & 0 & 0 \\ 1 & 0 & 1 \end{pmatrix} + S \begin{pmatrix} 0 & 0 & 2 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ & + 2S \begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 2 \\ 1 & 0 & 0 \end{pmatrix} + 2S \begin{pmatrix} 0 & 0 & 1 \\ 1 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix} \\ & + S \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{pmatrix} + S \begin{pmatrix} 1 & 0 & 1 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + S \begin{pmatrix} 1 & 0 & 1 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} + S \begin{pmatrix} 1 & 0 & 2 \\ 0 & 0 & 0 \\ 1 & 1 & 0 \end{pmatrix} \\ & + 2S \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix} + 2S \begin{pmatrix} 0 & 0 & 1 \\ 2 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix} + 2S \begin{pmatrix} 0 & 0 & 2 \\ 2 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}. \end{aligned}$$

$$(85) \quad \begin{aligned} S \begin{pmatrix} 0 & 3 \\ 1 & 1 \end{pmatrix} * S \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} = & S \begin{pmatrix} 2 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} + S \begin{pmatrix} 2 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + S \begin{pmatrix} 2 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 1 & 0 \end{pmatrix} + S \begin{pmatrix} 2 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{pmatrix} \\ & + 2S \begin{pmatrix} 1 & 0 & 1 \\ 2 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} + 2S \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix} \\ & + S \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} + S \begin{pmatrix} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + S \begin{pmatrix} 0 & 1 & 1 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{pmatrix} + S \begin{pmatrix} 0 & 2 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 1 \end{pmatrix} \\ & + 2S \begin{pmatrix} 0 & 0 & 1 \\ 1 & 2 & 1 \\ 0 & 0 & 0 \end{pmatrix} + 2S \begin{pmatrix} 0 & 1 & 0 \\ 1 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix} + 2S \begin{pmatrix} 0 & 2 & 0 \\ 0 & 0 & 2 \\ 1 & 0 & 0 \end{pmatrix}. \end{aligned}$$

$$\begin{aligned}
(86) \quad S \begin{pmatrix} 3 & 0 \\ 1 & 1 \end{pmatrix} * S \begin{pmatrix} 1 & 2 \\ 1 & 1 \end{pmatrix} = & S \begin{pmatrix} 0 & 0 & 2 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} + S \begin{pmatrix} 0 & 0 & 2 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{pmatrix} + S \begin{pmatrix} 1 & 0 & 2 \\ 0 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix} + S \begin{pmatrix} 0 & 0 & 2 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\
& + 2S \begin{pmatrix} 1 & 0 & 1 \\ 0 & 0 & 2 \\ 0 & 1 & 0 \end{pmatrix} + 2S \begin{pmatrix} 0 & 0 & 1 \\ 1 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix} \\
& + S \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{pmatrix} + S \begin{pmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} + S \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + S \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 1 \end{pmatrix} \\
& + 2S \begin{pmatrix} 1 & 0 & 0 \\ 1 & 2 & 1 \\ 0 & 0 & 0 \end{pmatrix} + 2S \begin{pmatrix} 0 & 1 & 0 \\ 2 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix} + 2S \begin{pmatrix} 0 & 2 & 0 \\ 2 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}.
\end{aligned}$$

$$\begin{aligned}
(87) \quad S \begin{pmatrix} 3 & 0 \\ 1 & 1 \end{pmatrix} * S \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} = & S \begin{pmatrix} 2 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} + S \begin{pmatrix} 2 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + S \begin{pmatrix} 2 & 1 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 1 \end{pmatrix} + S \begin{pmatrix} 2 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{pmatrix} \\
& + 2S \begin{pmatrix} 1 & 1 & 0 \\ 2 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} + 2S \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix} \\
& + S \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{pmatrix} + S \begin{pmatrix} 1 & 0 & 1 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + S \begin{pmatrix} 1 & 0 & 1 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} + S \begin{pmatrix} 2 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix} \\
& + 2S \begin{pmatrix} 0 & 0 & 1 \\ 2 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix} + 2S \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix} + 2S \begin{pmatrix} 2 & 0 & 0 \\ 0 & 0 & 2 \\ 0 & 1 & 0 \end{pmatrix}.
\end{aligned}$$

If one denotes by h the commutative image of S , one easily checks that

$$\begin{aligned}
(88) \quad h \begin{pmatrix} 3 \\ 1 \end{pmatrix} h \begin{pmatrix} 0 \\ 1 \end{pmatrix} * h \begin{pmatrix} 2 \\ 1 \end{pmatrix} h \begin{pmatrix} 1 \\ 1 \end{pmatrix} = & 2h \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} h \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} h \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + h \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix} h \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} h \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} + h \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix} h \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} h \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \\
& + 2h \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} h \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} h \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} + 2h \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} h \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} h \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \\
& + 2h \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} h \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} h \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + h \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} h \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} h \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} + h \begin{pmatrix} 2 \\ 0 \\ 0 \end{pmatrix} h \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} h \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} \\
& + 4h \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} h \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix} h \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + 2h \begin{pmatrix} 2 \\ 0 \\ 0 \end{pmatrix} h \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix} h \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}.
\end{aligned}$$

6. THE MANTACI-REUTENAUER ALGEBRA

6.1. Monochromatic complete functions. Let $\mathbf{e}_i$ be the canonical basis of $\mathbb{N}^\ell$. For $n \geq 1$, let

$$(89) \quad S_n^{(i)} = S_{n \cdot \mathbf{e}_i} \in \mathbf{Sym}^{(\ell)},$$

be the *monochromatic complete symmetric functions*.

Proposition 6.1. *The $S_n^{(\ell)}$ generate a Hopf subalgebra $\text{MR}^{(\ell)}$ of $\text{Sym}^{(\ell)}$, which is isomorphic to the Mantaci-Reutenauer descent algebra of the wreath products $\mathfrak{S}_n^{(\ell)} = (\mathbb{Z}/\ell\mathbb{Z}) \wr \mathfrak{S}_n$ if $C = \mathbb{Z}/\ell\mathbb{Z}$.*

Proof – $\text{MR}^{(\ell)}$ is obviously stable by the product and coproduct coming from $\text{Sym}^{(\ell)}$, hence is a Hopf subalgebra of $\text{Sym}^{(\ell)}$.

As a Hopf algebra, it is clearly isomorphic to the Mantaci-Reutenauer algebra, having the same number of independent generators in each degree, and behaving in the same way (divided powers) under the coproduct. The isomorphism for the internal product comes from the splitting formula, which gives explicitly $S^{(I,u)} * S^{(J,v)}$. ■

Since $\text{MR}^{(\ell)}$ has ℓ generators in each degree, its dimensions are given by

$$(90) \quad \frac{1}{1 - \sum_{k \geq 1} \ell t^k} = 1 + \ell \sum_{n \geq 1} (\ell + 1)^{n-1} t^n.$$

The bases of $\text{MR}^{(\ell)}$ are labelled by colored compositions, that is, pairs formed by a composition and a color vector of the same length:

$$(91) \quad (I, u) = ((i_1, \dots, i_m), (u_1, \dots, u_m)).$$

6.2. Primitive elements of $\text{MR}^{(\ell)}$. $\text{MR}^{(\ell)}$ being a subalgebra of a graded connected cocommutative Hopf algebra, is itself a graded connected cocommutative Hopf algebra, so that, thanks to the Cartier-Milnor-Moore theorem, it is the universal enveloping algebra of $L^{(\ell)}$:

$$(92) \quad \text{MR}^{(\ell)} = U(L^{(\ell)}),$$

where $L^{(\ell)}$ is the Lie algebra of its primitive elements. The following property is obtained in the same way as Theorem 4.2.

Theorem 6.2. *As a graded Lie algebra, the primitive Lie algebra $L^{(\ell)}$ of $\text{MR}^{(\ell)}$ is free over a set indexed by colored compositions.*

For example, with $\ell = 2$, the generating series of the dimensions of $L^{(\ell)}$ is

$$(93) \quad 1 + 2t + 3t^2 + 8t^3 + 18t^4 + 48t^5 + 116t^6 + 312t^7 + \dots$$

With $\ell = 3$, one finds

$$(94) \quad 1 + 3t + 6t^2 + 20t^3 + 60t^4 + 204t^5 + 670t^6 + 2340t^7 + \dots$$

More generally, the dimension of $L_n^{(\ell)}$ is given by the Witt polynomials

$$(95) \quad q_n(\ell) := \begin{cases} \ell & \text{if } n = 1, \\ \frac{1}{n} \sum_{d|n} \mu(d) (\ell + 1)^{n/d} & \text{if } n > 1, \end{cases}$$

so that, for $n \geq 2$, the dimension of $L_n^{(\ell)}$ coincide with those of a free Lie algebra on $\ell + 1$ generators of degree 1.

6.3. Duality. The duality is easily worked out by means of the appropriate Cauchy kernel. The generating function of the complete functions is

$$(96) \quad \sigma_{\mathbf{x}}^{\text{MR}}(\mathbf{A}) := 1 + \sum_{j=0}^{\ell-1} \sum_{n \geq 1} S_n^{(j)} \cdot (x^{(j)})^n,$$

and the Cauchy kernel is as usual

$$(97) \quad K^{\text{MR}}(\mathbb{X}, \mathbf{A}) = \prod_{i \geq 1}^{\rightarrow} \sigma_{\mathbf{x}_i}^{\text{MR}}(\mathbf{A}) = \sum_{(I,u)} M_{(I,u)}(\mathbb{X}) S^{(I,u)}(\mathbf{A}),$$

where (I, u) runs over colored compositions $(I, u) = ((i_1, \dots, i_m), (u_1, \dots, u_m))$. The $M_{I,u}$ are called the *monochromatic monomial quasi-symmetric functions* and satisfy

$$(98) \quad M_{(I,u)}(\mathbb{X}) = \sum_{j_1 < \dots < j_m} (x_{j_1}^{(u_1)})^{i_1} \dots (x_{j_m}^{(u_m)})^{i_m}.$$

Proposition 6.3. *The $M_{(I,u)}$ span a subalgebra of $\mathbb{C}[X]$ which can be identified with the graded dual of $\text{MR}^{(\ell)}$ through the pairing*

$$(99) \quad \langle M_{(I,u)}, S^{(J,v)} \rangle = \delta_{I,J} \delta_{u,v},$$

where δ is the Kronecker symbol.

Note that this algebra can also be obtained by imposing the relations

$$(100) \quad x_i^{(p)} x_i^{(q)} = 0, \text{ for } p \neq q$$

on the variables of $QSym^{(\ell)}$.

Baumann and Hohlweg [2] have another construction of the dual of $\text{MR}^{(\ell)}$ (implicitly defined in [35], Lemma 11).

7. LEVEL ℓ PARKING QUASI-SYMMETRIC FUNCTIONS

7.1. Usual parking functions.

7.1.1. The combinatorial objects. Recall that a *parking function* on $[n] = \{1, 2, \dots, n\}$ is a word $\mathbf{a} = a_1 a_2 \dots a_n$ of length n on $[n]$ whose nondecreasing rearrangement $\mathbf{a}^\uparrow = a'_1 a'_2 \dots a'_n$ satisfies $a'_i \leq i$ for all i . Let PF_n be the set of such words. It is well-known that $|\text{PF}_n| = (n+1)^{n-1}$.

One says that $\mathbf{a}$ has a *breakpoint* at b if $|\{a_i \leq b\}| = b$. The set of all breakpoints of $\mathbf{a}$ is denoted by $BP(\mathbf{a})$. Then, $\mathbf{a} \in \text{PF}_n$ is said to be *prime* if $BP(\mathbf{a}) = \{n\}$ (see [42]).

Let $\text{PPF}_n \subset \text{PF}_n$ be the set of prime parking functions on $[n]$. It can easily be shown that $|\text{PPF}_n| = (n-1)^{n-1}$ (see [43]).

Finally, one says that $\mathbf{a}$ has a *match* at b if $|\{a_i < b\}| = b-1$ and $|\{a_i \leq b\}| \geq b$. The set of all matches of $\mathbf{a}$ is denoted by $\text{Ma}(\mathbf{a})$.

7.1.2. *Algebraic structure on parking functions.* The algebra **PQSym** of parking functions [29, 31] is very similar to the algebra **FQSym** of permutations.

Since parking functions are closed under the shifted shuffle, one defines a product on the vector space with basis $(\mathbf{F}_{\mathbf{a}})$ by

$$(101) \quad \mathbf{F}_{\mathbf{a}'} \mathbf{F}_{\mathbf{a}''} = \sum_{\mathbf{a} \in \mathbf{a}' \uplus \mathbf{a}'' [k]} \mathbf{F}_{\mathbf{a}}.$$

The coproduct on this basis is given by the parkization algorithm [29]: for $w = w_1 w_2 \cdots w_n$ on $\{1, 2, \dots\}$, let us define

$$(102) \quad d(w) := \min\{i \mid \#\{w_j \leq i\} < i\}.$$

If $d(w) = n + 1$, then w is a parking function and the algorithm terminates, returning w . Otherwise, let w' be the word obtained by decrementing all the elements of w greater than $d(w)$. Then $\text{Park}(w) := \text{Park}(w')$. Since w' is smaller than w in the lexicographic order, the algorithm terminates and always returns a parking function.

For example, let $w = (3, 5, 1, 1, 11, 8, 8, 2)$. Then $d(w) = 6$ and the word $w' = (3, 5, 1, 1, 10, 7, 7, 2)$. Then $d(w') = 6$ and $w'' = (3, 5, 1, 1, 9, 6, 6, 2)$. Finally, $d(w'') = 8$ and $w''' = (3, 5, 1, 1, 8, 6, 6, 2)$, that is a parking function. Thus, $\text{Park}(w) = (3, 5, 1, 1, 8, 6, 6, 2)$.

We then have

$$(103) \quad \Delta \mathbf{F}_{\mathbf{a}} = \sum_{\substack{u, v \\ u \cdot v = \mathbf{a}}} \mathbf{F}_{\text{Park}(u)} \otimes \mathbf{F}_{\text{Park}(v)}.$$

7.1.3. *Duality.* Let $\mathbf{G}_{\mathbf{a}} = \mathbf{F}_{\mathbf{a}}^* \in \mathbf{PQSym}^*$ be the dual basis of $(\mathbf{F}_{\mathbf{a}})$. If $\langle \cdot, \cdot \rangle$ denotes the duality bracket, the product on $\mathbf{PQSym}^*$ is given by

$$(104) \quad \mathbf{G}_{\mathbf{a}'} \mathbf{G}_{\mathbf{a}''} = \sum_{\mathbf{a}} \langle \mathbf{G}_{\mathbf{a}'} \otimes \mathbf{G}_{\mathbf{a}''}, \Delta \mathbf{F}_{\mathbf{a}} \rangle \mathbf{G}_{\mathbf{a}} = \sum_{\mathbf{a} \in \mathbf{a}' * \mathbf{a}''} \mathbf{G}_{\mathbf{a}},$$

where the *convolution* $\mathbf{a}' * \mathbf{a}''$ of two parking functions is defined as

$$(105) \quad \mathbf{a}' * \mathbf{a}'' = \sum_{\substack{u, v; \mathbf{a} = u \cdot v, \\ \text{Park}(u) = \mathbf{a}', \text{Park}(v) = \mathbf{a}''}} \mathbf{a}.$$

By duality, one easily gets the formula for the coproduct of $\mathbf{G}_{\mathbf{a}}$ as

$$(106) \quad \Delta \mathbf{G}_{\mathbf{a}} := \sum_{u, v; \mathbf{a} \in u \uplus v} \mathbf{G}_u \otimes \mathbf{G}_v.$$

7.2. **Colored parking functions.** Let ℓ be an integer, representing the number of allowed colors. A *colored parking function* of level ℓ and size n is a pair composed of a parking function of length n and a word on $[\ell]$ of length n .

Since there is no restriction on the coloring, it is obvious that there are $\ell^n (n+1)^{n-1}$ colored parking functions of level ℓ and size n .

With two colors, one finds the sequence $a_i = 2(2i+2)^{i-1}$, known as A097629 in [41]:

$$(107) \quad 1 + 2t + 12t^2 + 128t^3 + 2000t^4 + 41472t^5 + 1075648t^6 + \dots$$

Since the convolution of two parking functions contains only parking functions, one easily builds as in [29] an algebra $\mathbf{PQSym}^{(\ell)}$ on colored parking functions:

$$(108) \quad \mathbf{G}_{(\mathbf{a}', u')} \mathbf{G}_{(\mathbf{a}'', u'')} = \sum_{\mathbf{a} \in \mathbf{a}' * \mathbf{a}''} \mathbf{G}_{(\mathbf{a}, u' \cdot u'')}.$$

We can define a coproduct using sums of alphabets: again, we only need a total order on A to define the colored parkization, so that taking two isomorphic copies A' and A'' of A , we define $\mathbf{A}' \oplus \mathbf{A}''$ as $(A' \oplus A'') \times C$, where $A' \oplus A''$ denotes the ordered sum. Assuming furthermore that $\mathbf{A}'$ and $\mathbf{A}''$ commute, we identify $f(\mathbf{A}')g(\mathbf{A}'')$ with $f \otimes g$ and define a coproduct by:

$$(109) \quad \Delta f(\mathbf{A}) = f(\mathbf{A}' \oplus \mathbf{A}'').$$

By construction, this is an algebra morphism from $\mathbf{PQSym}^{(\ell)}$ to $\mathbf{PQSym}^{(\ell)} \otimes \mathbf{PQSym}^{(\ell)}$, so that

Theorem 7.1. $\mathbf{PQSym}^{(\ell)}$ is a graded connected bialgebra, hence a Hopf algebra. More precisely, the coproduct can be computed in the following way:

$$(110) \quad \Delta \mathbf{G}_{(\mathbf{a}, u)} = \sum_{\substack{(\mathbf{a}', \mathbf{a}'', u', u'') \\ (\mathbf{a}, u) \in (\mathbf{a}', u') \uplus (\mathbf{a}'', u'')}} \mathbf{G}_{(\mathbf{a}', u')} \otimes \mathbf{G}_{(\mathbf{a}'', u'')}.$$

Proof – Straightforward from the previous definitions. ■

For example,

$$(111) \quad \Delta \mathbf{G}_{(41142, 22115)} = 1 \otimes \mathbf{G}_{(41142, 22115)} + \mathbf{G}_{(112, 215)} \otimes \mathbf{G}_{(11, 21)} + \mathbf{G}_{(41142, 22115)} \otimes 1.$$

7.3. Duality. Let $\mathbf{F}_{(\mathbf{a}, u)} = \mathbf{G}_{(\mathbf{a}, u)}^* \in \mathbf{PQSym}^{(\ell)*}$ be the dual basis of $(\mathbf{G}_{\mathbf{a}})$. If $\langle , \rangle$ denotes the duality bracket, the product on $\mathbf{PQSym}^{(\ell)*}$ is given by

$$(112) \quad \mathbf{F}_{(\mathbf{a}', u')} \mathbf{F}_{(\mathbf{a}'', u'')} = \sum_{(\mathbf{a}, u) \in (\mathbf{a}', u') \uplus (\mathbf{a}'', u'')} \mathbf{F}_{(\mathbf{a}, u)},$$

where the *shifted shuffle* of two colored parking functions is such that colors follow their letters.

Using the duality bracket once more, one easily gets the formula for the coproduct of $\mathbf{F}_{(\mathbf{a}, u)}$ as

$$(113) \quad \Delta \mathbf{F}_{(\mathbf{a}, u)} = \sum_{\substack{(p', u'), (p'', u'') \\ p' p'' = \mathbf{a}; u' u'' = u}} \mathbf{F}_{(\text{Park}(p'), u')} \otimes \mathbf{F}_{(\text{Park}(p''), u'')}.$$

7.4. Algebraic structure of $\mathbf{PQSym}^{(\ell)}$. Recall that a word w over $\mathbb{N}^*$ is *connected* if it cannot be written as a shifted concatenation $w = u \bullet v$, and *anti-connected* if its mirror image $\overline{w}$ is connected.

We denote by $\mathcal{CP}$ the set of connected parking functions, and by $p_n := |\mathcal{CP}_n|$ the number of such parking functions of size n . For later reference, we recall that the generating series of p_n is Sequence A122708 of [41]:

$$\begin{aligned} p(t) &:= \sum_{n \geq 1} p_n t^n = 1 - \left(\sum_{n \geq 0} (n+1)^{n-1} t^n \right)^{-1} \\ &= t + 2t^2 + 11t^3 + 92t^4 + 1014t^5 + 13795t^6 + 223061t^7 + \dots \end{aligned}$$

Let the *connected colored parking functions* be the $(\mathbf{a}, u)$ with $\mathbf{a}$ connected and u arbitrary. Their generating series is given by $p(\ell t)$.

From [31], we immediately get

Proposition 7.2. $\mathbf{PQSym}^{(\ell)}$ is free over the set $\mathbf{F}_{\sigma, u}$ (or $\mathbf{G}_{\sigma, u}$), where (σ, u) is connected.

For example, one gets the following generating series for the algebraic generators (connected parking functions with $\ell = 2$):

$$(114) \quad 2t + 8t^2 + 88t^3 + 1472t^4 + 32448t^5 + 882880t^6 + 28551808t^7 \dots$$

7.5. Primitive elements of $\mathbf{PQSym}^{(\ell)}$. Let $\mathcal{L}'^{(\ell)}$ be the primitive Lie algebra of $\mathbf{PQSym}^{(\ell)}$. Since Δ is not cocommutative, $\mathbf{PQSym}^{(\ell)}$ cannot be the universal enveloping algebra of $\mathcal{L}'^{(\ell)}$. But since it is cofree, it is, according to [23], the universal enveloping dipterous algebra of its primitive part $\mathcal{L}'^{(\ell)}$.

Let $\mathbf{G}^{\mathbf{a}, u}$ be the multiplicative basis defined by $\mathbf{G}^{\mathbf{a}, u} = \mathbf{G}_{\mathbf{a}_1, u_1} \cdots \mathbf{G}_{\mathbf{a}_r, u_r}$ where $(\mathbf{a}, u) = (\mathbf{a}_1, u_1) \bullet \cdots \bullet (\mathbf{a}_r, u_r)$ is the unique maximal factorization of $(\mathbf{a}, u) \in \mathfrak{S}_n \times C^n$ into connected colored parking functions.

Proposition 7.3. Let $\mathbf{V}_{\mathbf{a}, u}$ be the adjoint basis of $\mathbf{G}^{\mathbf{a}, u}$. Then, the family $(\mathbf{V}_{\alpha, u})_{\alpha \in \mathcal{CP}}$ is a basis of $\mathcal{L}'^{(\ell)}$. In particular, we have $\dim \mathcal{L}'^{(\ell)}_n = \ell^n p_n$. Moreover, $\mathcal{L}'^{(\ell)}$ is free.

Proof – The first part of the statement follows from [4]. The second part comes from the fact that $\mathbf{PQSym}^{(\ell)}$ is bidendriform (Theorem 7.4 below). $\blacksquare$

For example, since $\mathcal{L}'^{(\ell)}$ is free, the generating series of the degree of its generators is (with $\ell = 2$):

$$\begin{aligned} (115) \quad 1 - \prod_{n \geq 1} (1 - t^n)^{\ell^n p_n} &= 1 - (1 - t)^2 (1 - t^2)^8 (1 - t^3)^{88} \dots \\ &= 2t + 7t^2 + 72t^3 + 1276t^4 + 28944t^5 + 805288t^6 \\ &\quad + 26462232t^7 + \dots \end{aligned}$$

and the Hilbert series of the universal enveloping algebra of $\mathbf{PQSym}^{(\ell)}$ (its domain of cocommutativity) is, again with $\ell = 2$:

$$\begin{aligned} (116) \quad \prod_{n \geq 1} (1 - t^n)^{-\ell^n c_n} &= 1 + 2t + 11t^2 + 108t^3 + 1713t^4 + 36470t^5 + 969919t^6 \\ &\quad + 30847464t^7 + \dots \end{aligned}$$

7.6. Bidendriform and tridendriform structure.

Theorem 7.4. *The algebra $\mathbf{PQSym}^{(\ell)}$ has a structure of bidendriform bialgebra, hence is free as a Hopf algebra and as a dendriform algebra, cofree, self-dual, and its primitive Lie algebra is free.*

Moreover, the totally primitive elements of $\mathbf{PQSym}^{(\ell)}$ are the totally primitive elements of $\mathbf{PQSym}$ with any coloring.

Proof – It has been done in [31] in the case of $\mathbf{FQSym}$. But since the dendriform and codendriform structure do not involve the color alphabet C , the property is true in this case as well.

Also, colors do not play any role in determining whether a given element is (totally) primitive or not, so the last statement holds. $\blacksquare$

For example, the dendriform generators of $\mathbf{PQSym}$ have as degree generating series

$$(117) \quad \sum_i \text{dgp}_i t^i = t + t^2 + 7t^3 + 66t^4 + 786t^5 + 11\,378t^6 + 189\,391t^7 + \dots$$

so that the dendriform generators of $\mathbf{PQSym}^{(2)}$ have as degree generating series 2^idgp_i :

$$(118) \quad 2t + 4t^2 + 56t^3 + 1\,056t^4 + 25\,152t^5 + 721\,792t^6 + 24\,242\,048t^7 + \dots$$

7.6.1. Tridendriform structure.

Conjecture 7.5. *As in the case of $\mathbf{PQSym}$, we conjecture that $\mathbf{PQSym}^{(\ell)}$ is a free dendriform trialgebra.*

Note that there cannot be any relations, even tridendriform relations, among the elements $\mathbf{F}_{1,c}$ where $c \in C$, so that $\mathbf{PQSym}^{(\ell)}$ contains the free tridendriform algebra $\mathfrak{TD}^{(\ell)}$ on ℓ generators.

Recall that, if $\mathbf{PQSym}^{(\ell)}$ is free as a tridendriform algebra, its generators have as generating series

$$(119) \quad \text{TD} := \frac{\text{PQ} - 1}{2\text{PQ}^2 - \text{PQ}},$$

where PQ is the generating series of $\mathbf{PQSym}^{(\ell)}$.

For example, the tridendriform generators of $\mathbf{PQSym}$ have as degree generating series

$$(120) \quad \sum_i \text{tgp}_i t^i = t + 5t^3 + 50t^4 + 634t^5 + 9\,475t^6 + 163\,843t^7 + \dots$$

so that the tridendriform generators of $\mathbf{PQSym}^{(2)}$ have as degree generating series 2^itgp_i :

$$(121) \quad 2t + 40t^3 + 800t^4 + 20\,288t^5 + 606\,400t^6 + 20\,971\,904t^7 + \dots$$

8. TYPE B ALGEBRAS

8.1. Parking functions of type B . In [38], Reiner defined non-crossing partitions of type B by analogy to the classical case. In our context, he defined the level 2 case. This allowed him to derive, by analogy with a simple representation theoretical result, a definition of parking functions of type B as the words on $[n]$ of size n .

We shall build another set of words, also enumerated by n^n that sheds light on the relation between type- A and type- B parking functions and provides a natural Hopf algebra structure on the latter.

First, fix two colors $0 < 1$. We say that a pair of words $(\mathbf{a}, u)$ composed of a parking function and a binary colored word is a *level 2 parking function* if

- the only elements of $\mathbf{a}$ that can have color 1 are the matches of $\mathbf{a}$.
- for all element of $\mathbf{a}$ of color 1, all letters equal to it and to its left also have color 1,
- all elements of $\mathbf{a}$ have at least once the color 0.

For example, there are 27 level 2 parking functions of size 3: there are the 16 usual ones all with full color 0, and the eleven new elements

$$(122) \quad \begin{aligned} & (111, 100), (111, 110), (112, 100), (121, 100), (211, 010), \\ & (113, 100), (131, 100), (311, 010), (122, 010), (212, 100), (221, 100). \end{aligned}$$

The first time the first rule applies is with $n = 4$, where one has to discard the words $(1122, 0010)$ and $(1122, 1010)$ since 2 is not a match of 1122. On the other hand, both words $(1133, 0010)$ and $(1133, 1010)$ are B_4 -parking functions since 1 and 3 are matches of 1133.

Theorem 8.1. *The restriction of $\mathbf{PQSym}^{(2)}$ to the $\mathbf{F}$ elements indexed by level 2 parking functions is a subalgebra of $\mathbf{PQSym}^{(2)}$. The restriction of $\mathbf{PQSym}^{(2)}$ to the $\mathbf{G}$ elements indexed by level 2 parking functions is a subcoalgebra of $\mathbf{PQSym}^{(2)}$.*

Proof – The shifted shuffle of two $\mathbf{F}$ elements indexed by level 2 parking functions only consists in level 2 parking functions since the definition only involves matches (preserved by shifted shuffle) and positions of colors 0 and 1 on a given letter inside a word, also preserved by shifted shuffle. The same property holds for the coproduct on the $\mathbf{G}$ side: a match on either side of the tensor product comes from a match of the original word and all equal letters go to the same side of the tensor product in the same order. ■

8.2. Non-crossing partitions of type B . Remark that in the level 1 case, non-crossing partitions are in bijection with non-decreasing parking functions. To extend this correspondence to type B , let us start with a non-decreasing parking function $\mathbf{b}$ (with no color). We factor it into the maximal shifted concatenation of prime non-decreasing parking functions, and we choose a color, here 0 or 1, for each factor. We obtain in this way $\binom{2n}{n}$ words π , which can be identified with *type B non-crossing partitions*.

Let

$$(123) \quad \mathbf{P}^\pi = \sum_{\mathbf{a}^\uparrow = \pi} \mathbf{F}_{\mathbf{a}}$$

where $w^\uparrow$ denotes the nondecreasing rearrangement of the letters of w . Then,

Theorem 8.2. *The $\mathbf{P}^\pi$, where π runs over the above set of non-decreasing signed parking functions, form the basis of a cocommutative Hopf subalgebra $\mathbf{NCPQSym}^{(2)}$ of $\mathbf{PQSym}^{(2)}$.*

Proof – The subalgebra part comes from the fact that the shifted shuffle does not mix prime factors. The coalgebra part selects pieces of each factor, hence satisfies that each letter of a (new) factor has identical color. The cocommutativity part comes from the fact that all rearrangements of a given word are considered at the same time. ■

All this can be extended to higher levels in a straightforward way: allow each prime non-decreasing parking function to choose any color among ℓ and use the factorization as above. Since non-decreasing parking functions are in bijection with Dyck words, the choice can be described as: each block of a Dyck word with no return-to-zero, chooses one color among ℓ . In this version, the generating series is obviously given by

$$(124) \quad \frac{1}{1 - \ell \frac{1 - \sqrt{1 - 4t}}{2}}.$$

For $\ell = 3$, we obtain Sequence A007854 of [41].

9. COLORED ANALOGS OF PLANAR BINARY TREES: $\mathbf{PBT}^{(\ell)}$

9.1. Definition of $\mathbf{PBT}^{(\ell)}$. In the case with one color, the Hopf algebra $\mathbf{PBT}$ of Planar binary trees initially defined by Loday and Ronco [22] can be embedded in $\mathbf{FQSym}$ in the following way [12, 13]:

$$(125) \quad \mathbf{P}_T = \sum_{\sigma; \text{shape}(\mathcal{P}(\sigma))=T} \mathbf{F}_\sigma,$$

where $\mathcal{P}$ is a simple algorithm: it is the well-known *binary search tree insertion*, such as presented, for example, by Knuth in [18].

In the colored case, the definition is almost the same:

$$(126) \quad \mathbf{P}_{T,u} = \sum_{(\sigma,u); \text{shape}(\mathcal{P}(\sigma))=T} \mathbf{F}_{(\sigma,u)},$$

where u is a color word whose length is equal to the number of leaves of T . Note that this algebra (without its realization on words) has been previously studied by Maria Ronco [40].

Given the definitions, the generating series of the dimensions of $\mathbf{PBT}^{(\ell)}$ is

$$(127) \quad 1 + \ell t + 2\ell^2 t^2 + 5\ell^3 t^3 + 14\ell^4 t^4 + \dots$$

that is, the generating series of Catalan numbers multiplied by ℓ^n .

9.2. Algebraic structure of $\mathbf{PBT}^{(\ell)}$. Since $\mathbf{PBT}$ is generated by the trees with no right branch (starting from the root), the same holds in $\mathbf{PBT}^{(\ell)}$:

Proposition 9.1. *$\mathbf{PBT}^{(\ell)}$ is free over the set $\mathbf{P}_{T,u}$, where T is a tree with no right branch.*

For example, the generating series of the algebraic generators of $\mathbf{PBT}^{(\ell)}$ is

$$(128) \quad \ell t + \ell^2 t^2 + 2\ell^3 t^3 + 5\ell^4 t^4 + 14\ell^5 t^5 + \dots$$

that is, the generating series of shifted Catalan numbers C_{n-1} multiplied by ℓ^n .

9.2.1. Primitive elements of $\mathbf{PBT}^{(\ell)}$. Let $\mathcal{L}^{(\ell)}$ be the primitive Lie algebra of the algebra $\mathbf{PBT}^{(\ell)}$. Since Δ is not cocommutative, $\mathbf{PBT}^{(\ell)}$ cannot be the universal enveloping algebra of $\mathcal{L}^{(\ell)}$. But since it is cofree, it is, according to [23], the universal enveloping dipterous algebra of its primitive part $\mathcal{L}^{(\ell)}$.

Using the same arguments as in the case of $\mathbf{FQSym}^{(\ell)}$, one then proves

Proposition 9.2. *The Lie algebra $\mathcal{L}^{(\ell)}$ is free. Moreover*

$$(129) \quad \dim \mathcal{L}_n^{(\ell)} = \ell^n C_{n-1},$$

For example, since $\mathcal{L}^{(\ell)}$ is free, the generating series of the degree of its generators is (with $\ell = 2$):

$$(130) \quad \begin{aligned} 1 - \prod_{n \geq 1} (1 - t^n)^{\ell^n C_{n-1}} &= 1 - (1 - t)^2 (1 - t^2)^4 (1 - t^3)^{16} (1 - t^4)^{80} \dots \\ &= 2t + 3t^2 + 8t^3 + 46t^4 + 252t^5 \\ &\quad + 1558t^6 + 9800t^7 + \dots \end{aligned}$$

and the Hilbert series of the universal enveloping algebra of $\mathbf{PBT}^{(\ell)}$ (its domain of cocommutativity) is, again with $\ell = 2$,

$$(131) \quad \begin{aligned} \prod_{n \geq 1} (1 - t^n)^{-\ell^n C_{n-1}} &= 1 + 2t + 7t^2 + 28t^3 + 139t^4 + 762t^5 + 4549t^6 \\ &\quad + 28464t^7 + \dots \end{aligned}$$

9.2.2. Dendriform structure of $\mathbf{PBT}^{(\ell)}$. Recall that $\mathbf{PBT}$ is the free dendriform algebra on one generator. Since colors do not play any role in determining if a given element is (totally) primitive or not, the same holds for $\mathbf{PBT}^{(\ell)}$:

Proposition 9.3. *The algebra $\mathbf{PBT}^{(\ell)}$ is the free dendriform algebra on ℓ generators. It has also the structure of bidendriform bialgebra.*

Note 9.4. It is also possible to define a colored analog of $\mathbf{CQSym}$ the Catalan Quasi-symmetric algebra, but the natural definition leads to a non-cocommutative algebra, hence not sharing the basic property of $\mathbf{CQSym}$ itself.

10. EXAMPLES

10.1. Multigraded coinvariants and colored Klyachko idempotents. One of the very first applications of the theory of noncommutative symmetric functions was to provide an explanation for the following coincidence. On the one hand, consider the representation of $\mathfrak{S}_n$ in the coinvariant algebra

$$(132) \quad \mathcal{H}_n = \mathbb{C}[x_1, \dots, x_n]_{\mathfrak{S}_n} = \mathbb{C}[x_1, \dots, x_n] / \mathcal{J},$$

where $\mathcal{J}$ is the ideal generated by symmetric polynomials without constant term. It is known [25] that the graded Frobenius characteristic of the action of $\mathfrak{S}_n$ on $\mathcal{H}_n$ is

$$(133) \quad \begin{aligned} \text{ch}_q(\mathcal{H}_n) &= \sum_k q^k \text{ch}(\mathcal{H}_n)^{(k)} \\ &= (q)_n h_n \left(\frac{X}{1-q} \right) = \sum_{I \models n} q^{\text{maj}(I)} r_I(X), \end{aligned}$$

where $r_I(X)$ are the ribbon Schur functions.

On the other hand, Klyachko [17] introduced a remarkable Lie idempotent in $\mathbb{C} \mathfrak{S}_n$

$$(134) \quad \kappa_n = \sum_{\sigma \in \mathfrak{S}_n} \zeta^{\text{maj}(\sigma)} \sigma$$

where ζ is a primitive n -th root of unity.

In terms of noncommutative symmetric functions, both expressions can be interpreted as specializations of

$$(135) \quad K_n(q) := (q)_n S_n \left(\frac{A}{1-q} \right) = \sum_{I \models n} q^{\text{maj}(I)} R_I.$$

This is the graded noncommutative characteristic of an action of $H_n(0)$ on coinvariants. This is a projective module, hence also an $\mathfrak{S}_n$ -module, and taking the commutative image, ($A = X$), one obtains the characteristic of $\mathfrak{S}_n$. But $K_n(q)$ can also be interpreted as an element of the descent algebra of $\mathfrak{S}_n$. A simple computation shows that for $q = \zeta$, a primitive n -th root of unity, $K_n(\zeta)$ is a primitive element of **Sym** of commutative image p_n/n , hence is a Lie idempotent (see [20]). Actually, $K_n(q)$ is the noncommutative Hall-Littlewood function $\tilde{H}_{1^n}(A; q)$, and this specialization property is a special case of a noncommutative version [11] of the classical property [11, 21, 25].

There is a similar phenomenon here. Let $q_1, \dots, q_n$ be independent variables, and consider the noncommutative symmetric function

$$(136) \quad \mathbf{K}_n(A; q_1, \dots, q_n) := \sum_{I \models n} \mathbf{q}^{\text{MAJ}(I)} R_I,$$

where

$$(137) \quad \mathbf{q}^{\text{MAJ}(i_1, \dots, i_r)} := (q_1 \dots q_{i_1})^r (q_{i_1+1} \dots q_{i_1+i_2})^{r-1} \dots (q_{i_1+\dots+i_{r-2}} \dots q_{i_1+\dots+i_{r-1}}).$$

For example,

$$(138) \quad \mathbf{K}_3 = R_3 + q_1 q_2 R_{21} + q_1 R_{12} + q_1^2 q_2 R_{111}.$$

$$(139) \quad \begin{aligned} \mathbf{K}_4 = & R_4 + q_1 q_2 q_3 R_{31} + q_1 q_2 R_{22} + q_1^2 q_2^2 q_3 R_{211} \\ & + q_1 R_{13} + q_1^2 q_2 q_3 R_{121} + q_1^2 q_2 R_{112} + q_1^3 q_2^2 q_3 R_{1111}. \end{aligned}$$

Its commutative image is the multigraded characteristic of $\mathcal{H}_n$ with respect to the *partition degree* (cf. [3]).

One may also regard $\mathbf{K}_n$ as an element of

$$(140) \quad \mathbf{Sym}_n^{(\mathbb{Z})} \subset \mathbf{FQSym}_n^{(\mathbb{Z})}$$

by means of the identification

$$(141) \quad R_I(A) q_1^{\alpha_1} \dots q_n^{\alpha_n} = \sum_{\text{Des}(\sigma)=I} \mathbf{G}_{\sigma, \alpha_1 \dots \alpha_n}$$

and writing it as

$$(142) \quad \mathbf{K}_n(A; q_1, \dots, q_n) = \sum_{I \models n} R_I(A) \mathbf{q}^{\text{MAJ}(I)}$$

Note that this element lives in the (descent) algebra of the extended affine Weyl group of type A

$$(143) \quad \widehat{\mathfrak{S}}_n = \mathbb{Z}^n \ltimes \mathfrak{S}_n = P \ltimes \mathfrak{S}_n,$$

where P is the weight lattice. One can also interpret it as an element of the usual affine Weyl group

$$(144) \quad \widetilde{\mathfrak{S}}_n = Q \ltimes \mathfrak{S}_n$$

where Q is the root lattice

$$(145) \quad Q = \{\alpha \in P \mid \sum_{i=1}^n \alpha_i = 0\}.$$

This amounts to imposing the relation

$$(146) \quad q_1 \dots q_n = 1,$$

which replaces the root of unity condition $q^n = 1$ in the one-parameter case.

It has been proved by McNamara and Reutenauer [28] that under condition (146), $\mathbf{K}_n(A; q_1, \dots, q_n)$ was a Lie idempotent in $\mathbb{C} \widetilde{\mathfrak{S}}_n$. Within the formalism of the present paper, this can be seen as follows: the authors of [28] introduce a twisted product on $\mathcal{A}_n = \mathbb{C}(x_1, \dots, x_n)[\mathfrak{S}_n]$ by the formula

$$(147) \quad f(\mathbf{x})\sigma \cdot g(\mathbf{x})\tau = f(\mathbf{x})\sigma[g(\mathbf{x})]\sigma\tau,$$

where permutations act on functions as automorphisms, *i.e.*, $\sigma(x_i) = x_{\sigma(i)}$, and in particular, on monomials by $\sigma[\mathbf{x}^c] = \mathbf{x}^{c\sigma^{-1}}$. Hence,

$$(148) \quad (\sigma\mathbf{x}^c) \cdot (\tau\mathbf{x}^d) = \sigma\tau\mathbf{x}^{c\tau+d},$$

which is the same as Formula (29) so that $\mathcal{A}_n$ can be identified with the homogeneous component of degree n of $\mathbf{FQSym}^{(\mathbb{Z})}$, by setting

$$(149) \quad \sigma\mathbf{x}^c \equiv \mathbf{G}_\sigma\mathbf{x}^c \equiv \mathbf{G}_{\sigma,c}.$$

McNamara and Reutenauer then introduce the formal series in $\mathbf{FQSym}^{(\mathbb{Z})}$

$$(150) \quad \Theta(\mathbf{x}) = \sum_{n \geq 0} \sum_{\sigma \in \mathfrak{S}_n} \frac{\prod_{j \in \text{Des}(\sigma)} x_{\sigma(1)} \cdots x_{\sigma(j)}}{\prod_{i=1}^n (1 - x_{\sigma(1)} \cdots x_{\sigma(i)})} \sigma$$

which, applying (148), and under the identification (149), can be rewritten as

$$(151) \quad \Theta(\mathbf{x}) = \prod_{l \geq 0} \sum_{n \geq 0} \mathbf{G}_{\text{id}_n, l^n} = \prod_{l \geq 0} \sigma_1(A^{(l)}).$$

Indeed, introducing the new variables

$$(152) \quad y_j = x_1 x_2 \cdots x_j,$$

and applying (148) we can write

$$(153) \quad \begin{aligned} \Theta(\mathbf{x}) &= \sum_{n \geq 0} \sum_{\sigma \in \mathfrak{S}_n} \mathbf{G}_\sigma \frac{\prod_{d \in \text{Des}(\sigma)} y_d}{(1 - y_1)(1 - y_2) \cdots (1 - y_n)} \\ &= \sum_{n \geq 0} \mathbf{K}_n(y_1, \dots, y_{n-1}) \frac{1}{((y))_n}, \end{aligned}$$

where $((y))_n = (1 - y_1)(1 - y_2) \cdots (1 - y_n)$ and

$$(154) \quad \mathbf{K}_n(y_1, \dots, y_{n-1}) = \sum_{\sigma \in \mathfrak{S}_n} \mathbf{G}_\sigma \prod_{d \in \text{Des}(\sigma)} y_d$$

is the twisted version of the multiparameter Klyachko element introduced in [20, (11)]. By Moebius inversion over the lattice of compositions of n , we have

$$(155) \quad \mathbf{K}_n(y_1, \dots, y_{n-1}) = \sum_{I \models n} R_I \cdot \prod_{d \in \text{Des}(I)} y_d$$

$$(156) \quad \mathbf{K}_n(y_1, \dots, y_{n-1}) \frac{1}{((y))_n} = \sum_{J \models n} S^J \cdot \frac{1}{1 - y_n} \prod_{d \in \text{Des}(J)} \frac{y_d}{1 - y_d},$$

which implies the expression

$$(157) \quad \Theta(\mathbf{x}) = \prod_{l \geq 0} \left(\sum_{n \geq 0} S_n y_n^l \right).$$

Each factor of the right-hand side is grouplike (for the coproduct of $\mathbf{FQSym}^{(l)}$):

$$(158) \quad \Delta \sum_{n \geq 0} S_n y_n^l = \sum_{n \geq 0} \sum_{i+j=n} S_i y_i^l \otimes S_j y_j^l = \sum_{i \geq 0} S_i y_i \otimes \sum_{j \geq 0} S_j y_j^l$$

so that also

$$(159) \quad \Delta \Theta(\mathbf{x}) = \Theta(\mathbf{x}) \otimes \Theta(\mathbf{x}).$$

Extracting the term of degree n , we find

$$(160) \quad \Delta \mathbf{K}_n = \sum_{i+j=n} \left(\mathbf{K}_i \frac{1}{((y))_i} \otimes \mathbf{K}_j \frac{1}{((y))_j} \right) ((y))_n$$

so that if we send y_n to 1, all terms vanish except the extreme ones, and we get a primitive element. This is the main result of [28].

10.2. A formula of Raney. Raney [37] gave a combinatorial interpretation of the coefficients of the unique solution $g(t) \in \mathbb{Q}[Y, Z][[t]]$ of the functional equation

$$(161) \quad g(t) = t \sum_{k=1}^{\ell} y_k e^{z_k g(t)},$$

with $g(t) = \sum_{n \geq 0} \frac{g_n}{n!} t^{n+1}$. This defining equation is of the form

$$(162) \quad g(t) = t\phi(g(t)),$$

with $\phi(u) = \sum_{k=1}^{\ell} y_k e^{z_k u}$. Hence, the coefficient g_n of t^{n+1} in $g(t)$ is

$$(163) \quad \begin{aligned} g_n &= \frac{1}{n+1} [u^n] \phi(u)^{n+1} \\ &= \frac{1}{n+1} \sum_{\substack{n_1 + \dots + n_{\ell} = n+1 \\ q_1 + \dots + q_{\ell} = n}} \binom{n+1}{n_1, \dots, n_{\ell}} \binom{n}{q_1, \dots, q_{\ell}} y_1^{n_1} \dots y_{\ell}^{n_{\ell}} (n_1 z_1)^{q_1} \dots (n_{\ell} z_{\ell})^{q_{\ell}}. \end{aligned}$$

Thus, $g_n \in \mathbb{N}[Y, Z]$. Its combinatorial interpretation can be mechanically derived by means of a colored version of the noncommutative Lagrange inversion formula as formulated in [33, 8]. Consider the functional equation

$$(164) \quad g = \sum_{k,n} b_k S_n^{(k)} g^n,$$

where $S_n^{(k)} = S_n(A^{(k)})$ is a colored complete function and b_k are noncommuting letters. We can set

$$(165) \quad S_n = \sum_k b_k S_n^{(k)},$$

so that (164) can be rewritten as

$$(166) \quad g = \sum_{n \geq 0} S_n g^n,$$

and the solution of [33] reads

$$(167) \quad \begin{aligned} g &= S^0 + S^{10} + (S^{200} + S^{110}) + \dots \\ &= \sum_{\pi \in \text{NDPF}} S^{\text{Ev}(\pi).0}, \end{aligned}$$

where NDPF is the set of nondecreasing parking functions. Note that $S^0 = \sum b_k$ is *a priori* different from 1, and does not commute with the other S^i . Each term $S^{\text{Ev}(\pi).0}$ represents an ordered tree T in Polish notation, so that, for example

$$(168) \quad \begin{array}{c} \circ \\ \swarrow \quad \searrow \\ \circ \qquad \circ \\ \qquad \quad | \\ \qquad \quad \circ \end{array}$$

is S^{2010} .

Replacing each S^i by $\sum_{k=1}^{\ell} b_k S_i^{(k)}$, the expression $S^{\text{Ev}(\pi),0}$ becomes a sum over all ℓ -colorings of the tree T , so that, one recovers the combinatorial interpretation of Raney (proved in a different way by Foata [5]): let $\mathbf{n} = (n_1, \dots, n_{\ell})$ and $\mathbf{q} = (q_1, \dots, q_{\ell}) \in \mathbb{N}^{\ell}$ and let $B(\mathbf{n}, \mathbf{q})$ be the set of ℓ -colored trees on $n = |\mathbf{n}|$ vertices with n_k vertices of color k and such that the sum of the arities of vertices of color k is q_k . Then

$$(169) \quad g = \sum_{n \geq 0} \frac{1}{n!} \sum |B(\mathbf{n}, \mathbf{q})| \mathbf{y}^{\mathbf{n}} \mathbf{z}^{\mathbf{q}}.$$

REFERENCES

- [1] P. BIANE, *Parking functions of types A and B*, Electronic J. Combin. **9** (2002), # 7.
- [2] P. BAUMANN and C. HOHLWEG, *A Solomon-type epimorphism for Mantaci-Reutenauer's algebra of a wreath product $G \wr \mathfrak{S}_n$* , Trans. Amer. Math. Soc. **360** (2008), 1475–1538.
- [3] F. CASELLI, *Diagonal invariants and the refined multimahonian distribution*, preprint, arXiv:math.CO/0805.2860.
- [4] G. DUCHAMP, F. HIVERT, and J.-Y. THIBON, *Noncommutative symmetric functions VI: free quasi-symmetric functions and related algebras*, Internat. J. Alg. Comput. **12** (2002), 671–717.
- [5] D. FOATA, *La série génératrice exponentielle dans les problèmes d'énumération*, Montréal, Presses de l'Univ. de Montréal, 1974.
- [6] L. FOISSY, *Bidendriform bialgebras, trees, and free quasi-symmetric functions*, J. Pure Appl. Algebra **209** (2007), no. 2, 439–459.
- [7] I.M. GELFAND, D. KROB, A. LASCoux, B. LECLERC, V. S. RETAKH, and J.-Y. THIBON, *Noncommutative symmetric functions*, Adv. in Math. **112** (1995), 218–348.
- [8] I. GESSEL, *Noncommutative Generalization and q -analog of the Lagrange Inversion Formula*, Trans. Amer. Math. Soc. **257** (1980), no. 2, 455–482.
- [9] I. GESSEL, *Multipartite P -partitions and inner products of skew Schur functions*, [in “Combinatorics and algebra”, C. Greene, Ed.], Contemporary Mathematics, **34** (1984), 289–301.
- [10] I. GESSEL, *Enumerative Applications of Symmetric Functions*, Sémin. Lothar. Combin., **17a** (1987), 17p. (electronic).
- [11] F. HIVERT, *Hecke algebras, difference operators, and quasi-symmetric functions*, Adv. Math., **155** (2000), no. 2, 181–238.
- [12] F. HIVERT, J.-C. NOVELLI, and J.-Y. THIBON, *Un analogue du monoïde plaxique pour les arbres binaires de recherche*, C. R. Acad. Sci. Paris Sér I Math. **332** (2002), 577–580.
- [13] F. HIVERT, J.-C. NOVELLI, and J.-Y. THIBON, *The algebra of binary search trees*, Theo. Comp. Sci. **339** (1) (2005), 129–165.
- [14] F. HIVERT, J.-C. NOVELLI, and J.-Y. THIBON, *Yang-Baxter bases of 0 -Hecke algebras and representation theory of 0 -Araki-Koike-Shoji algebras*, Adv. in Math. **205** (2006), 504–548.
- [15] F. HIVERT and N. THIÉRY, *MuPAD-Combinat, an open-source package for research in algebraic combinatorics*, Sémin. Lothar. Combin. **51** (2004), 70p. (electronic).
- [16] V. G. KAC, *Infinite-dimensional Lie algebras*, Cambridge Univ. Press, 1991.
- [17] A.A. KLYACHKO, *Lie elements in the tensor algebra*, Siberian Math. J. **15** (1974), 1296–1304.
- [18] D. E. KNUTH, *The art of computer programming, vol.3: Sorting and searching*, (Addison-Wesley, 1973).
- [19] A. G. KONHEIM and B. WEISS, *An occupancy discipline and applications*, SIAM J. Applied Math. **14** (1966), 1266–1274.
- [20] D. KROB, B. LECLERC, and J.-Y. THIBON, *Noncommutative symmetric functions II: Transformations of alphabets*, Internal J. Alg. Comput. **7** (1997), 181–264.
- [21] A. LASCoux, B. LECLERC and J.-Y. THIBON, *Green polynomials and Hall-Littlewood functions at roots of unity*, Europ. Jour. Comb. **15** (1994), 173–180.

- [22] J.-L. LODAY and M.O. RONCO, *Hopf algebra of the planar binary trees*, Adv. Math. **139** (1998) n. 2, 293–309.
- [23] J.-L. LODAY and M.O. RONCO, *Algèbres de Hopf colibres*, C. R. Acad. Sci. Paris Sér I Math. **337** (2003), 153–158.
- [24] J.-L. LODAY, *Scindement d’associativité et algèbres de Hopf*, Actes des Journées Mathématiques à la Mémoire de Jean Leray, Sémin. Congr. Soc. Math. France 9 (2004), 155–172.
- [25] I.G. MACDONALD, *Symmetric functions and Hall polynomials*, 2nd ed., Oxford University Press, 1995.
- [26] P.A. MCMAHON, *Combinatory analysis*, Cambridge University Press, 1915, 1916; Chelsea reprint, 1960.
- [27] R. MANTACI and C. REUTENAUER, *A generalization of Solomon’s descent algebra for hyperoctahedral groups and wreath products*, Comm. Algebra **23** (1995), 27–56.
- [28] P. MCNAMARA and C. REUTENAUER, *P-Partitions and a Multi-Parameter Klyachko Idempotent*, Elec. Jour. Comb. **11**(2) (2005), #R21.
- [29] J.-C. NOVELLI and J.-Y. THIBON, *A Hopf Algebra of Parking Functions*, Proc. FPSAC’04 Conf., Vancouver.
- [30] J.-C. NOVELLI and J.-Y. THIBON, *Construction of dendriform trialgebras*, C.-R. Acad. Sci., Paris Sr. I **342** (2006), 365–369.
- [31] J.-C. NOVELLI and J.-Y. THIBON, *Hopf algebras and dendriform structures arising from parking functions*, Funda. Math. **193** (2007), 189–241.
- [32] J.-C. NOVELLI and J.-Y. THIBON, *Free quasi-symmetric functions of arbitrary level*, preprint, math.CO/0405597.
- [33] J.-C. NOVELLI and J.-Y. THIBON, *Noncommutative Symmetric Functions and Lagrange Inversion*, Intern. Jour. Alg. and Comput, to appear.
- [34] F. PATRAS, *L’algèbre des descentes d’une bigèbre graduée*, J. Algebra **170** (1994), 547–566.
- [35] S. POIRIER, *Cycle type and descent set in wreath products*, Disc. Math., **180** (1998), 315–343.
- [36] S. POIRIER and C. REUTENAUER, *Algèbre de Hopf des tableaux*, Ann. Sci. Math. Québec **19** (1995), 79–90.
- [37] G. RANEY, *A formal solution of $\sum_{i=1}^{\infty} A_i e^{B_i X} = X$* , Can. J. Math. **16** (1964), 755–762.
- [38] V. REINER, *Non-crossing partitions for classical reflection groups*, Preprint available at <ftp://s6.math.umn.edu/pub/papers/reiner/>
- [39] C. REUTENAUER, *Free Lie algebras*, Oxford University Press, 1993.
- [40] M.O. RONCO, *Primitive elements in a free dendriform Hopf algebra*, Contemp. Maths. **267** (2000) 245–264.
- [41] N.J.A. SLOANE, The On-Line Encyclopedia of Integer Sequences, <http://www.research.att.com/~njas/sequences/>
- [42] R. P. STANLEY, *Parking functions and noncrossing partitions*, Electronic J. Combin. **4** (1997), # 2.
- [43] R. P. STANLEY, *Enumerative combinatorics*, vol. 2, Cambridge University Press, 1999.

UNIVERSITÉ PARIS-EST, INSTITUT GASPARD MONGE, 5 BOULEVARD DESCARTES, CHAMPS-SUR-MARNE, 77454 MARNE-LA-VALLÉE CEDEX 2, FRANCE

E-mail address, Jean-Christophe Novelli: novelli@univ-mlv.fr

E-mail address, Jean-Yves Thibon: jyt@univ-mlv.fr